

Finiteness conditions and infinite matrix rings ^{*}

Ángel del Río and Juan Jacobo Simón [†]

November 19, 2004

Abstract

For a unital ring R , $\text{RCFM}_\alpha(R)$ denotes the ring of row and column finite matrices over R indexed by α . We give necessary and sufficient structural conditions on $\text{RCFM}_\alpha(R)$ which are equivalent to R being, respectively, Quasi-Frobenius, left artinian, and left noetherian.

In this paper R denotes an (associative and unital) ring and α is an infinite set. We use the following notation, where “matrices” means “matrices indexed by α with entries in R ”:

$$\begin{aligned} A = \text{RFM}_\alpha(R) &= \text{Ring of row finite matrices,} \\ B = \text{RCFM}_\alpha(R) &= \text{Ring of row and column finite matrices,} \\ B_0 = \text{FM}_\alpha(R) &= \text{Ring of finite matrices.} \end{aligned} \tag{1}$$

At a first sight it might seem that the rings A and B are too big to reflect properties of the ring R , and still more unexpected that A or B could encode finiteness conditions of R . However, already in [6, 12] it is shown that the ring A reflects some finiteness conditions of R . There is a long tradition in the study of the ring theoretical properties of the ring A (among others see [2, 6, 9, 12]). Recently several authors have shown interest in the study of the ring B (see e.g. [5, 7, 10]). In this paper we study the properties of the ring B under the assumption that R satisfies some finiteness condition (quasi-Frobenius, artinian, noetherian).

The relationship between a ring R and A comes essentially from the adjoint pair $\text{Hom}_R(F, -) : R\text{-mod} \rightleftharpoons A\text{-mod} : F \otimes_A -$, where F is a free left R -module of rank $|\alpha|$ and A is canonically identified with $\text{End}_R(F)$. An interesting exception may be found in the computation of the Jacobson radical (see [13]) where the amount of matrix manipulation exceeds adjunction techniques. If one wants to relate the rings R and B one can also use the adjoint pair $\text{Hom}_R(F, -) : R\text{-mod} \rightleftharpoons B\text{-mod} : F \otimes_B -$. The difference in the performance of the adjoint pairs for A and B relies on the fact that while A is in the image of $\text{Hom}_R(F, -)$, namely $A \simeq \text{End}_R(F)$, this is not the case for B . However, we may still use some “adjoint-like” techniques to relate some special objects in the category of R -modules and some special matrices or ideals of B and keeping in mind that B is the ring of continuous endomorphisms of F in a certain topology [11].

We start with some notation. If $i, j \in \alpha$ then e_{ij} denotes the element of B_0 having 1 in the (i, j) -th entry and zeroes elsewhere, and if $a \in A$ then $a(i, j)$ denotes the (i, j) -th entry of a . Set $e_i = e_{ii} = e_{\{i\}}$. If F is a subset of α then set $e_F = \sum_{i \in F} e_i$. A careful study of the arithmetic of the rings A , B and B_0 and their idempotents e_F for F finite leads to the following lemma.

^{*}1991 Mathematics Subject Classification 16S50 (16P20,16P40)

[†]Partially supported by the D.G.I. of Spain and Fundación Séneca of Murcia

Lemma 1 1. B_0 is a two sided ideal of B and a right ideal of A .

2. For every $x \in B_0$ there is $e = e^2 \in B_0$ such that $x = ex = xe$.

3. The map $\rho : A \rightarrow \text{End}_{(B_0 B_0)}$ that associates $a \in A$ with the endomorphism ρ_a of B_0 given by $\rho_a(x) = xa$ is a ring isomorphism.

We need the following module theoretical lemma (see [4, Exercise 18.17]).

Lemma 2 A left R -module M is quasi-injective if and only if $f(M) \subseteq M$ for every endomorphism f of the injective hull of M .

Recall that R is said to be quasi-Frobenius (QF) if R is right and left artinian, and there exists a duality between the categories of finitely generated left and right R -modules. Further R is said to be quasi-continuous if $e(R) \subseteq R$ for every idempotent endomorphism e of the injective hull of ${}_R R$.

In [6] it is shown that R is QF if and only if A is left self injective. In contrast, the ring B cannot be even left or right quasi-continuous [7] for any ring R . Our first result shows how to compute the injective hull of ${}_B B$ if R is QF. Notice that as a consequence of the next theorem any of the equivalent conditions 2-6 hold for some infinite set α if and only if they hold for every infinite set.

Theorem 3 The following conditions are equivalent for a ring R and an infinite set α :

1. R is quasi-Frobenius.

2. $\text{RFM}_\alpha(R)$ is left self-injective.

3. $\text{FM}_\alpha(R)$ is an injective object in the category $\text{FM}_\alpha(R) - \text{mod}$ consisting of those $\text{FM}_\alpha(R)$ -modules M such that $\text{FM}_\alpha(R)M = M$.

4. $\text{FM}_\alpha(R)$ is quasi-injective as a left $\text{RCFM}_\alpha(R)$ -module.

5. $\text{RFM}_\alpha(R)$ is injective as a left $\text{RCFM}_\alpha(R)$ -module.

6. $\text{RFM}_\alpha(R)$ is the injective hull of the regular left $\text{RCFM}_\alpha(R)$ -module $\text{RCFM}_\alpha(R)$.

Proof. Let A, B and B_0 as in (1) and let E denote the injective hull of ${}_B B$.

The equivalence between 1 and 2 was proved in [6].

The equivalence between 5 and 6 follows from the fact that B_0 is essential in ${}_B A$ and hence $E = E({}_B A) = E({}_B B_0)$.

3 implies 4 follows from the equality $B_0 = BB_0$.

Identifying A with $\text{End}_R(R^{(\alpha)})$ one can consider $R^{(\alpha)}$ as a $R - B_0$ -bimodule and it is well known that the functor $F = B_0 \text{Hom}_R(R^{(\alpha)}, -) : R - \text{mod} \rightarrow B_0 - \text{mod}$ is an equivalence of categories such that $F(R^{(\alpha)}) = B_0$. (Alternatively one can prove this by using the results of [1] or [3].) Now the equivalence between 1 and 3 follows from the fact R is quasi-Frobenius if and only if ${}_R R^{(\alpha)}$ is injective; this is a direct consequence of Theorems 24.18 and 24.20 of [8].

Now we prove that 4 implies 6. Assume that ${}_B B_0$ is quasi-injective. We have already seen that $A \subseteq E$. Let $e \in E$ and consider the map $f : B_0 \rightarrow E$ given by $f(x) = xe$. By assumption and Lemmas 1 and 2 there is an $a \in A$ such that $xe = xa$ for every $x \in B_0$. We claim that $e = a$ and this completes the proof. Indeed, if $e \neq a$ then $B(e - a) \neq 0$ and since B_0 is essential in ${}_B E$ there is $0 \neq x \in B_0 \cap B(e - a)$. Thus there is $x_0 \in B_0$ and $b \in B$ such that $x = x_0 x = b(e - a)$ and hence $x = x_0 x = x_0 b(e - a) = 0$, a contradiction.

Finally we prove that 5 implies 3. Assume that ${}_B A$ is injective and let $f : N \rightarrow B_0$ be a homomorphism of B_0 -modules with N a submodule of B_0 . By using $B_0 N = N$ and $B_0 = B_0 B = B B_0$ one deduces that f is a homomorphism of B -modules and hence f extends to an endomorphism g of ${}_B A$. If $x \in B_0$ then there is $e \in B_0$ such that $x = ex$ and hence $g(x) = eg(x) \in B_0$. Therefore g restricts to an endomorphism of ${}_B B_0$ which extends f . This shows that ${}_B B_0$ is quasi-injective by Lemma 2. ■

For every $i \in \alpha$, consider the map $\pi_i : B \rightarrow R^{(\alpha)}$ that associates to $a \in B$, the i -th row of a . Let I be a left ideal of B . Since $\pi_j(a) = \pi_i(e_{ij} \cdot a)$, $\pi_i(I)$ does not depend on i . Write $\pi(I) = \pi_i(I)$ for $i \in \alpha$ arbitrary. Clearly $\pi(I)$ is a submodule of ${}_R R^{(\alpha)}$. (There is an alternative definition of $\pi(I)$ for I a left ideal of B as $R^{(\alpha)} I$ where $R^{(\alpha)}$ is considered as a R - B -bimodule in the natural matricial way.) Thus if $a \in B$ then $\pi(Aa)$ is the submodule of $R^{(\alpha)}$ generated by the rows of a . If M is a submodule of ${}_R R^{(\alpha)}$ then $\iota(M) = \{a \in B : \pi(a) \in M, \text{ for every } i \in \alpha\}$ is a left ideal of B .

Definition 4 We say that a left ideal I of B is closed if for every $a \in B \setminus I$ there is $i \in I$ such that $\pi_i(a) \notin \pi(I)$.

Lemma 5 A left ideal I of B is closed if and only if $I = \iota(M)$ for some submodule M of ${}_R R^{(\alpha)}$. In particular the left annihilators $l_B(X)$ in B of subsets X of B and the left ideals of the form $B(J) = \{a \in B : a(i, j) \in J \text{ for every } i, j\}$ where J is a left ideal of R are closed left ideals of B .

Proof. Clearly $I \subseteq \iota(\pi(I))$ and I closed if and only if the equality holds. Further if M is a submodule of ${}_R R^{(\alpha)}$ then $\iota(\pi(\iota(M))) = \iota(M)$. This proves the first statement. The second is a consequence of the first because $B(J) = \iota(J^{(\alpha)})$ and if $X \subseteq B$ then $l_B(X) = \iota(M)$ where M is the annihilator in ${}_R R^{(\alpha)}$ of X . ■

Proposition 6 The following conditions are equivalent for a left ideal J of R :

1. $\text{RCFM}_\alpha(J) = \{a \in \text{RCFM}_\alpha(R) : a(i, j) \in J \text{ for every } i, j\}$ is a cyclic left ideal of $\text{RCFM}_\alpha(R)$.
2. $\text{RCFM}_\alpha(J)$ is a finitely generated left ideal of $\text{RCFM}_\alpha(R)$.
3. J is finitely generated left ideal of R .

Proof. Let A, B and B_0 as in (1). The equivalence between 1 and 2 follows from the fact that B has single basis number, i.e. ${}_B B \simeq {}_B B^n$ for every positive integer n . If ${}_B B(J)$ is generated by X then J is generated by the entries of the first row of the elements of X . This proves 2 implies 3.

Let x_1, \dots, x_n generate ${}_R J$. Since α is infinite there is a natural identification between $B(J)$, $\text{RCFM}_\alpha(M_{n,1}(K))$ and $\text{RCFM}_\alpha(M_{1,n}(R))$ that we are going to use without specific mention. Let $a \in B(J)$ be the following "diagonal" matrix (diagonal in $M_\alpha(M_{n,1}(J))$)

$$g = \begin{pmatrix} X & 0 & 0 & \cdots \\ 0 & X & 0 & \cdots \\ 0 & 0 & X & \cdots \\ \vdots & \vdots & \vdots & \ddots \end{pmatrix} \quad \text{with} \quad X = \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix}.$$

If $b \in B(J)$ then for every $x, y \in J$ there is $c(x, y) \in M_{1,n}(R)$ such that the (x, y) -th entry $b(x, y)$ of b is $c(x, y)X$ and we may take $c(x, y) = 0$ if $b(x, y) = 0$. Then $c = (c(x, y))_{x, y \in \alpha} \in B$ and $b = ca$. This proves that $B(J) = Ba$. ■

Remark 7 If RCFM is replaced by RFM in conditions 1 and 2 of Proposition 6 then they are still equivalent, since $\text{RFM}_\alpha(R)$ has single basis number. However they are not equivalent to 3. For example if J is the ideal generated by a subset X of R then $\text{RFM}_X(J)$ is the cyclic left ideal generated by the matrix having the elements of X in one column and zeroes in the remaining columns.

The Jacobson radical of R is denoted by $J(R)$. The ring R is said to be a Baer ring if any left (equivalently, right) annihilator in R of a subset of R is a direct summand.

Theorem 8 If α is an infinite set then a unital ring R is left artinian and if and only if $J(\text{RCFM}_\alpha(R))$ is cyclic as a left ideal of $\text{RCFM}_\alpha(R)$ and $\text{RCFM}_\alpha(R)/J(\text{RCFM}_\alpha(R))$ is a Baer ring. In this case $J(\text{RCFM}_\alpha(R)) = \text{RCFM}_\alpha(J(R))$.

Proof. Assume that R is left artinian. By Proposition 6, $\text{RCFM}_\alpha(J(R))$ is a cyclic left ideal of $\text{RCFM}_\alpha(R)$. By [7, Theorem 14], $\text{RCFM}_\alpha(R)/J(\text{RCFM}_\alpha(R))$ is a Baer ring, and $J(\text{RCFM}_\alpha(R)) = \text{RCFM}_\alpha(J(R))$; so that $J(\text{RCFM}_\alpha(R))$ is a cyclic left ideal of $\text{RCFM}_\alpha(R)$.

Conversely, by [7, Theorem 14], R is a perfect ring and $J(\text{RCFM}_\alpha(R)) = \text{RCFM}_\alpha(J(R))$. By hypothesis and Proposition 6, $J(R)$ is a finitely generated left ideal of R . Then by [4, Ex. 28.9] R is left artinian. ■

By Proposition 6 if every closed left ideal of B is finitely generated then R is left noetherian. Our next result proves the converse for α countable.

Theorem 9 The following conditions are equivalent for a unital ring R :

1. R is left noetherian.
2. Every closed left ideal of $\text{RCFM}_{\mathbb{N}}(R)$ is finitely generated.
3. Every closed left ideal of $\text{RCFM}_{\mathbb{N}}(R)$ is cyclic.

Proof. By Proposition 6 we only have to prove 1 implies 3.

Let $I = \iota(M)$ be a closed left ideal of $B = \text{RCFM}_{\mathbb{N}}(R)$. For every non negative integer n let $p_n : R^{(\mathbb{N})} \rightarrow R^{(\mathbb{N})}$ be the projection on the first n coordinates and let

$$M_n = M \cap \text{Im } p_n \quad \text{and} \quad K_n = M \cap \ker p_n.$$

Notice that

$$0 = M_0 \leq M_1 \leq M_2 \leq \dots \leq \bigcup_{n \geq 0} M_n = M = K_0 \geq K_1 \geq K_2 \geq \dots \geq \bigcap_{n \geq 0} K_n = 0.$$

If M is finitely generated, say $M = \langle x_1, \dots, x_k \rangle$ then I is generated by the matrix

$$a = \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_k \\ 0 \\ \vdots \end{pmatrix}.$$

(Notice that the rows of a are elements of $R^{(\mathbb{N})}$ and hence $a \in B$.) Thus assume that M is not finitely generated. This implies that $M \neq M_n$ for every $n \geq 0$.

If $n \geq 0$, then $p_n(M)$ is finitely generated and hence $p_n(M) = p_n(\langle m_1, \dots, m_k \rangle)$ for some $m_1, \dots, m_k \in M$. Since M is not finitely generated there is $m \in M \setminus \langle m_1, \dots, m_k \rangle$. On the other hand $p_n(m) = \sum_{i=1}^k r_i p_n(m_i)$ for some $r_i \in R$ and thus $0 \neq m - \sum_{i=1}^k r_i m_i \in K_n$.

The argument of the previous paragraph shows that $K_n \neq 0$ for every $n \geq 0$ and using this fact we construct recursively a sequence $(m_n, k_n, S_n)_{n \geq 0}$ as follows:

$$\begin{aligned} k_0 &= 0, \text{ the integer;} \\ S_0 &= 0, \text{ the trivial submodule of } M; \\ m_n &= \min\{m : M_m \cap K_{k_n} \neq 0\}; \\ k_{n+1} &= \min\{k : (M_{m_n} + \sum_{i=0}^n S_i) \cap K_k = 0\} \end{aligned}$$

and S_{n+1} is a finitely generated submodule of K_{k_n} such that

$$\left(M_{m_{n+1}} + \sum_{i=0}^n S_i \right) \cap K_{k_n} \subseteq S_{n+1} \subseteq K_{k_n} \quad \text{and} \quad (S_{n+1} + K_{k_{n+1}})/K_{k_{n+1}} = K_{k_n}/K_{k_{n+1}}. \quad (2)$$

The existence of m such that $M_m \cap K_{k_n} \neq 0$ is warranted by the fact that $K_{k_n} \neq 0$; the existence of k such that $(M_{m_n} + \sum_{i=0}^n S_i) \cap K_k = 0$ follows from the fact the $M_{m_n} + \sum_{i=0}^n S_i$ is finitely generated and hence it is embedded in K_k for some k ; finally the existence of S_{n+1} finitely generated satisfying (2) follows from the fact that $M_{m_{n+1}} \cap K_{k_n}$ and $K_{k_n}/K_{k_{n+1}}$ are finitely generated. Since $M_{m_n} \cap K_{k_n} \neq 0$, $M_{m_n} \cap K_{k_{n+1}} \neq 0$ and $M_{m_{n+1}} \cap K_{k_{n+1}} \neq 0$, $m_n < m_{n+1}$ and $k_n < k_{n+1}$ for every n , i.e. (m_n) and (k_n) are increasing sequences.

Claim: For every $n, r \geq 0$, $(M_{m_n} + \sum_{i=0}^n S_i) \cap K_{k_r} \subseteq \sum_{i \geq r+1}^{n+1} S_i$. In particular, $K_{k_r} \subseteq \sum_{i \geq r+1} S_i$ for every $r \leq 0$.

If $n - r < 0$ then $(M_{m_n} + \sum_{i=0}^n S_i) \cap K_{k_r} \subseteq (M_{m_n} + \sum_{i=0}^n S_i) \cap K_{k_{n+1}} = 0$ and the Claim is obvious, so we assume that $n - r \geq 0$ and argue by induction on $n - r$. If $n - r = 0$ then $(M_{m_n} + \sum_{i=0}^n S_i) \cap K_{k_r} = (M_{m_r} + \sum_{i=0}^r S_i) \cap K_{k_r} \subseteq S_{r+1}$ by construction. Assume that $n - r > 0$ and let $m \in (M_{m_n} + \sum_{i=0}^n S_i) \cap K_{k_r}$. Then there is $s_{r+1} \in S_{r+1}$ such that $n = m - s_{r+1} \in K_{k_{r+1}}$. Thus $n \in (M_{m_n} + \sum_{i=0}^n S_i) \cap K_{k_{r+1}}$, by induction hypothesis $n \in \sum_{i \geq r+2}^{n+1} S_i$ and so $m \in \sum_{i \geq r+1}^{n+1} S_i$. This proves the Claim.

For every n let X_n be a finite generating set of S_n and construct the matrix

$$a = \begin{pmatrix} X_1 \\ X_2 \\ X_3 \\ \vdots \end{pmatrix}$$

that is, the first rows of a are formed by the elements of X_1 , in some order, the next rows are formed by the elements of X_2 , etc. Since $S_n \subseteq K_{k_n}$ and (k_n) is a strictly increasing sequence, each column of a has only finitely many non zero entries, that is $a \in B$ and because I is a closed left ideal and every row of a belongs to $M = \pi(I)$ one deduces that $a \in I$.

Let u_n be the cardinality of X_n and $v_n = \sum_{i=1}^n u_n$. By the claim if $x \in K_r$ then $x = ya$ for some $y \in \ker p_{v_r}$. If $x \in I$ then there is a strictly increasing sequence $(r_n)_n$ such that $\pi_m(x) \in K_{k_n}$ for every $m \geq r_n$. Thus if $r_n \leq m < r_{n+1}$ then $\pi_m(x) = y_m a$ with $y_m \in \ker p_{v_r}$ and hence $x = ya$, where y is the row finite matrix defined by setting $\pi_m(y) = y_m$. Since $M = \sum_{n \geq 1} S_n$ is not finitely generated the sequence (v_n) is non decreasing and non bounded and this implies that $y \in B$. Thus $x \in Ba$ and this proves that $I = Ba$. ■

Notice that the proof of Theorem 9 does not apply if the index set is not countable. We do not know whether or not the closed ideals of $B_\alpha(R)$ are cyclic for R noetherian and α a non countable set. A consequence of Theorem 9 is the following.

Corollary 10 *If R is left noetherian then $\text{RCFM}_{\mathbb{N}}(R)$ is left coherent.*

Proof. Let I be a finitely generated left ideal of B . Since B has single basis number there is a surjective homomorphism $f : B \rightarrow I$ of left B -modules. Then $I = Ba$ for some $a \in B$ and hence $\text{Ker } f$ is the left annihilator of a . Thus $\text{Ker } f$ is a closed left ideal of B . By Theorem 9, $\text{Ker } f$ is finitely generated and we conclude that I is finitely presented. ■

We conjecture that the converse of Corollary 10 is false in general. However, we provide a partial converse in the next proposition.

Proposition 11 *Let α be an infinite set and R a unital ring. If $\text{RCFM}_\alpha(R)$ is left coherent then R is left coherent and satisfies acc on direct summands.*

Proof. Assume that $B = \text{RCFM}_\alpha(R)$ is left coherent. To prove that R is left coherent we show that if $a_1, \dots, a_r \in R$ then the kernel of the homomorphism $f : R^n \rightarrow R$ given by $f(x_1, \dots, x_n) = \sum_{i=1}^n x_i a_i$ is finitely generated. Let $J = \{j_1, \dots, j_n\}$ be a subset of α of cardinality n and $a \in B$ be given by

$$a(x, y) = \begin{cases} 1, & \text{if } x = y \notin J \\ a_i, & \text{if } x = j_i \text{ and } y = j_1 \\ 0, & \text{otherwise} \end{cases}$$

By the construction of a , the left annihilator $l_B(a)$ of a in B is formed by the element $b \in B$ such that $b(x, y) = 0$ if $y \notin J$ and $(b(x, j_1), \dots, b(x, j_n)) \in \text{ker } f$ for every $x \in \alpha$. By hypothesis there is $b \in B$ such that $Bb = l_B(a)$ and hence it is easy to see that $\text{ker } f$ is generated by the elements of the form $b(x) = (b(x, j_1), \dots, b(x, j_n))$ with $x \in \alpha$. Since $b \in B$, $b(x) = 0$ for almost all $x \in \alpha$ and hence $\text{ker } f$ is finitely generated as wanted.

Now we prove that R satisfies acc on direct summands. Otherwise R has an infinite countable set $\{f_1, f_2, \dots\}$ of non zero orthogonal idempotents. Let $J = \{j_1, j_2, \dots\}$ be an infinite countable subset of α and consider the matrix $a \in B$ given as follows:

$$a(x, y) = \begin{cases} 1, & \text{if } x = y \notin J \\ 1 - f_n, & \text{if } x = y = j_n \\ -f_{n+1}, & \text{if } x = j_{n+1} \text{ and } y = j_n \\ f_{n+1} - 1, & \text{if } x = j_{n+2} \text{ and } y = j_n \\ 0, & \text{otherwise} \end{cases}$$

By hypothesis $l_B(a) = Bb$ for some $b \in B$. Let $I = \sum_{n \in \mathbb{N}} Rf_n$ and K the left ideal of R generated by the entries in the j_1 -column of b . Notice that K is finitely generated while I is not. We are going to obtain a contradiction by showing that $I = K$.

To prove $I \subseteq K$, consider, for any $n \in \mathbb{N}$, the matrix $m_n \in B$ having f_n in the entries (j_1, j_i) for $1 \leq i \leq n$ and 0 in any other entry. Notice that $m_n a = 0$ and hence $m_n \in Bb$. Thus $f_n \in K$.

The reverse inclusion follows by showing that all the entries of b belong to I . (In fact the same holds for any element in $l_B(a)$). Fix $x \in \alpha$ and let $Y = \{y \in \alpha : b(x, y) \neq 0\}$. We have to prove that $b(x, y) \in I$ for every $y \in Y$. This is obvious if $Y = \emptyset$, so assume that this is not the case. Having in mind that $ba = 0$ one has that $Y \subseteq J$. Let n be the maximum positive integer such that $b(x, j_n) \neq 0$. We prove that $b(x, j_m) \in I$ for $1 \leq m \leq n$ by induction on

$m - n$. First $0 = (ba)(x, j_n) = b(x, j_n)(1 - f_n)$ and hence $b(x, j_n) \in Rf_n \subseteq I$. Second $0 = (ba)(x, j_{n-1}) = b(x, j_{n-1})(1 - f_{n-1}) - b(x, j_n)f_n$ and hence $b(x, j_{n-1}) \in Rf_{n-1} + Rf_n \subseteq I$. Finally, if $1 \leq m \leq n - 2$ then $0 = (ba)(x, j_m) = b(x, j_m)(1 - f_{j_m}) - b(x, j_{m+1})f_{m+1} + b(x, j_{m+1})(f_{m+1} - 1)$ and hence $b(x, j_m) \in I$, because by the induction hypothesis $b(x, j_{m+1}) \in I$. ■

Corollary 12 *If α is an infinite cardinal and R is a ring then R is semisimple artinian if and only if R is von Neumann regular and $\text{RCFM}_\alpha(R)$ is left coherent.*

Proof. This is a direct consequence of Proposition 11 and the fact that R is semisimple artinian if and only if R is von Neumann regular and satisfies acc on direct summands [8, Theorem 19.26 A]. ■

References

- [1] G.D. Abrams, *Morita equivalence for rings with local units*, Comm. Algebra 11 (1983) 801–837.
- [2] G. Abrams and J. Haefner, *Picard groups and infinite matrix rings*, Trans. Amer. Math. Soc. 350 (1998) 2737–2752.
- [3] P.N. Ánh and L. Márki, *Morita equivalence for rings without identity*, Tsukuba J. Math. 11 (1987) 1–16.
- [4] F.W. Anderson and K.R. Fuller, *Rings and categories of modules*, Springer-Verlag, 1974.
- [5] P. Ara, E. Pardo and F. Perera, *The structure of countably generated projective modules over regular rings*, J. Algebra 226 (2000) 161–190.
- [6] G.M. Brodskii, *Endomorphism rings of free modules*, Mat. Sb. 94(136) (1974), 226–242. Math. USSR Sbornik 23(2), (1974).
- [7] V. Camillo, F.J. Costa-Cano and J.J. Simón, *Relating properties of a ring and its ring of row and column finite matrices*, J. Algebra 224 (2001) 435–449.
- [8] C. Faith, *Algebra II. Ring Theory*, Springer-Verlag, 1976.
- [9] P. Menal, *On the endomorphism ring of a free module*, Publ. Sec. Mat. Univ. Autònoma Barcelona 27 (1983), no. 1, 141–154.
- [10] K.C. O’Meara, *The exchange property for row and column-finite matrix rings*, J. Algebra 268 (2003) no. 2, 744–749.
- [11] D. Ornstein, *Dual vector spaces*, Ann. Math. 69 (1959) 520–534.
- [12] B. L. Osofsky, *Some properties of rings reflected in endomorphism rings of free modules*, Cont. Math. **13** 1982, 179–181.
- [13] R. Ware and J. Zelmanowitz, *The Jacobson radical of the endomorphism ring of a projective module*, Proc. AMS, **26** (1970), 15–20.

Departamento de Matemáticas, Universidad de Murcia, 30100 Murcia, Spain.
adelrio@um.es, jsimon@um.es