

Birth of piglets after transferring of *in vitro*-produced embryos pre-matured with R-roscovitine

Pilar Coy, Raquel Romar, Salvador Ruiz, Sebastián Cánovas, Joaquín Gadea, Francisco García Vázquez and Carmen Matás

Departamento de Fisiología, Facultad de Veterinaria, Universidad de Murcia, Murcia 30071, Spain

Correspondence should be addressed to P Coy; Email: pcoy@um.es

Abstract

The objectives of this study were to evaluate: (1) the nuclear maturation, (2) the intracellular glutathione (GSH) content, (3) the normality of fertilization and (4) full development after transplantation of embryos derived from porcine oocytes pre-cultured with 50 $\mu\text{mol/l}$ roscovitine (an inhibitor of p34^{cdc2}/cyclin B kinase) for 22 h. After treatment with roscovitine, the nuclear configuration of oocytes (Hoechst staining) was comparable with those examined just after collection: the majority of oocytes were arrested at the germinal vesicle (GV) 1 stage (63.2%). Roscovitine-treated oocytes progressed through meiosis to the metaphase II stage in a conventional step-wise *in vitro* maturation (IVM) program for 44 h in a proportion similar to control ones (>85.0%). When roscovitine-treated oocytes and non-treated oocytes were matured for 44 h and then co-cultured with fresh spermatozoa for 18 h, no differences were observed in oocyte penetrability, proportion of monospermic penetration and male pronuclear formation (>87%). Roscovitine increased the GSH synthesis in oocytes at 22 h, whereas, after 44 h, roscovitine-treated oocytes had similar amounts of GSH to non-treated oocytes. Finally, surgical transfer of zygotes at 22–24 h post-insemination, derived from roscovitine-treated oocytes, resulted in one pregnancy with 12 piglets born; control non-treated zygotes resulted in one pregnancy and 10 piglets born. The full-term developmental ability of mammalian oocytes pre-cultured with roscovitine prior to IVM is thereby demonstrated. This validation is important before the introduction of roscovitine into routine procedures.

Reproduction (2005) **129** 747–755

Introduction

Efforts to characterize oocytes within antral follicles not yet destined to ovulate have been hampered by an inability to maintain meiotic arrest in oocytes after removal from the follicle (reviewed by Picton & Gosden 1999). Whereas *in vivo*, meiotically competent oocytes are arrested at the germinal vesicle (GV) stage by the follicular environment until the preovulatory surge of gonadotropins, *in vitro*, oocytes resume meiosis spontaneously after removal from the follicular environment. Attempts to develop *in vitro* culture systems to maintain bovine or porcine oocytes at the GV stage using different meiotic inhibitors have been reported, and the results have been partially successful. The increasing interest in such studies is supported by several potential applications: (1) obtaining oocytes from different places (farms) that could be later sent to the laboratory under meiotic arrest for further *in vitro* maturation (IVM) and fertilization (IVF) – as an example, this method is being applied for commercial ovum pick up-*in vitro* embryo production (OPU-IVP) production of cattle embryos (P Mermillod,

personal communication); (2) maintaining the oocyte's ability to synthesize and store molecules important for subsequent embryonic development during the inhibitory period, thus allowing *in vitro* production of competent oocytes (Kubelka *et al.* 2000, Mermillod *et al.* 2000, Motlik *et al.* 2000, Ponderato *et al.* 2001); (3) a improvement in the post-freezing survival rates of GV stage oocytes compared with those at the metaphase stage; (4) possible optimization of cell cycle synchrony and production of cells that respond more predictably to nuclear reprogramming (Gibbons *et al.* 2002); (5) prolonged and more flexible maturation schedules in the laboratory (Coy *et al.* 2004); (6) the development of *in vitro* models to study the effects of elevated temperature (heat stress) and other environmental factors on immature oocytes (Payton *et al.* 2004). Because of these applications, two-step culture systems (a pre-culture in a medium with inhibitor and a further culture in IVM medium) are proposed as useful tools in IVM programs (Motlik *et al.* 2000).

The resumption of meiosis is regulated by activation of the metaphase promoting factor (MPF; Wu *et al.* 1996,

Motlik *et al.* 1998, Sirard *et al.* 1998). MPF is a cell division kinase (cdk) whose activation involves the formation of a complex between p34^{cdc2} kinase (cdk1) and cyclin B. Once formed, this complex should be activated by dephosphorylation of the threonine 14 and tyrosine 15 residues of the p34 subunit. The most common approach to block MPF activation and, consequently, the resumption of meiosis, has been the employment of cdk inhibitors. Roscovitine actively competes for the ATP binding sites in the cdk1 subunit of the MPF, and was first used to prevent germinal vesicle breakdown (GVBD) in cattle and pig oocytes (Mermillod *et al.* 2000, Krischek & Meinelcke, 2001, Marchal *et al.* 2001, Ponderato *et al.* 2001, 2002). Specifically, the inhibitory effect of roscovitine on the cell cycle was discovered by Meijer (reviewed by Meijer & Raymond 2003). The results seem to demonstrate that it is more effective and produces less detrimental effects than other inhibitors such as cycloheximide (Faerge *et al.* 2001), butyrolactone I (Kitagawa *et al.* 1993, Meijer & Kim 1997, Kubelka *et al.* 2000, Fair *et al.* 2002) or 6-DMAP (Lonergan *et al.* 1997, Avery *et al.* 1998, Liu *et al.* 1998, Dode & Adona 2001).

Despite the numerous papers published in recent years, and even though studies in cattle have been more extensive than in pigs, no references are available showing the influence of pre-culture with roscovitine and further IVM on important parameters such as the nuclear stage during the maturation period, the intracellular glutathione (GSH) content of pre-cultured oocytes or male pronuclear formation after IVF of these gametes. Moreover, the birth of offspring has yet to be reported, and therefore also unreported is the competence to develop to term of embryos produced *in vitro* under the two-step IVM conditions. As a consequence, the overall objective of this experimental work was to assess the influence of roscovitine on the nuclear and cytoplasmic maturation of oocytes entering a two-step IVM system and to demonstrate its effectiveness to generate embryos with full-term developmental ability.

Materials and Methods

Culture media

Unless otherwise indicated, all chemicals and reagents were purchased from Sigma-Aldrich Química S.A. The R-enantiomer of roscovitine (2-(R)-(1-ethyl-2-hydroxyethylamino)-6-benzylamino-9-isopropylpurine; R-7772) was solubilized in dimethyl sulfoxide before freezing at -20°C as a 10 mmol/l stock. The medium used for oocyte maturation was NCSU-37 (Petters & Wells 1993) supplemented with 0.57 mmol/l cysteine, 1 mmol/l dibutyl cAMP, 5 $\mu\text{g/ml}$ insulin, 50 $\mu\text{mol/l}$ β -mercaptoethanol, 10 iu/ml equine chorionic gonadotropin (eCG) (Foligon; Intervet International B.V., Boxmeer, Holland), 10 iu/ml human CG (hCG) (Chorulon, Intervet International B.V.) and 10% porcine follicular fluid (v/v).

The basic medium used for IVF was essentially the same as that used by Rath *et al.* (1999). This medium, designated as TALP medium, consists of: 114.06 mmol/l NaCl, 3.2 mmol/l KCl, 8 mmol/l Ca-lactate \cdot 5H₂O, 0.5 mmol/l MgCl₂ \cdot 6H₂O, 0.35 mmol/l NaH₂PO₄, 25.07 mmol/l NaHCO₃, 10 ml/l Na-lactate, 1.1 mmol/l Na-pyruvate, 5 mmol/l glucose, 2 mmol/l caffeine, 3 mg/ml BSA (A-9647), 1 mg/ml polyvinyl alcohol (PVA) and 0.17 mmol/l kanamycin sulfate.

The embryo culture medium was NCSU-23 containing: 0.4% BSA (A-8022), 75 $\mu\text{g/ml}$ potassium penicillin G and 50 $\mu\text{g/ml}$ streptomycin sulphate (Machaty *et al.* 1998).

Oocyte collection and *in vitro* maturation

Within 30 min of slaughter, ovaries from prepubertal gilts were transported to the laboratory in saline containing 100 $\mu\text{g/ml}$ kanamycin sulfate at 38°C , washed once in 0.04% cetrimide solution and twice in saline. Oocyte-cumulus cell complexes (COCs) were collected from antral follicles (3–6 mm diameter) by slicing, washed twice with Dulbecco's PBS supplemented with 1 mg/ml PVA and twice more in maturation medium previously equilibrated for a minimum of 3 h at 38.5°C under 5% CO₂ in air. Only COCs with a complete and dense cumulus oophorus were used for the experiments (Coy *et al.* 2002). Groups of 50 COCs were cultured in 500 μl maturation medium for 22 h at 38.5°C under 5% CO₂ in air. After culture, oocytes were washed twice in fresh maturation medium without dibutyl cAMP, eCG and hCG and cultured for an additional 20–22 h (Funahashi & Day 1993). This IVM system is commonly employed in pigs for several years and is based upon the results from Funahashi *et al.* (1997).

Brilliant cresyl blue test

Immediately after collection, COCs were washed in PBS and exposed to brilliant cresyl blue (BCB; 13 $\mu\text{mol/l}$ in PBS) for 90 min at 38.5°C in a humidified air atmosphere. BCB staining determines the activity of glucose-6-phosphate dehydrogenase, an enzyme synthesized in growing oocytes but with less activity in grown oocytes (Roca *et al.* 1998, Rodríguez-González *et al.* 2002). After exposure to BCB, they were washed three times in PBS and observed under a stereomicroscope at $\times 20$ and classified into two groups, depending on their cytoplasm coloration: oocytes showing blue cytoplasm or grown oocytes (BCB+) and oocytes without blue coloration or growing oocytes (BCB-).

Hoechst staining

Oocytes were fixed for 15 min (2% glutaraldehyde in PBS), stained for 15 min (1% Hoechst 33342 in PBS) and finally washed in PBS containing 1 mg/ml polyvinylpyrrolidone and mounted on glass slides. Oocytes were examined under an epifluorescence microscope at $\times 200$ and

× 400 magnification and designated as GV-0, GV-I GV-II, GV-III, GV-IV (including diakinesis and prometaphase I), metaphase I (Met I), anaphase I (Ana I) and Met II stages according to the morphological criteria for characterization of meiotic stages by Funahashi *et al.* (1997).

GSH assay

The intracellular content of GSH was measured as described previously (Funahashi *et al.* 1995). Briefly, COCs were denuded by pipetting in PBS and washed three times in a buffer solution. Five microliters of buffer containing 30 oocytes per replicate and group were transferred to a 2 ml microfuge tube and 5 µl of 1.25 mol/l phosphoric acid added. Samples were frozen immediately (−20°C) and kept in the freezer until assayed. The GSH content in the oocytes was determined by the dithionitrobenzoic acid–glutathione disulphide (DTNB–GSSG) reductase recycling assay (Anderson 1985). Briefly, 700 µl of 0.33 mg/ml NADPH in 0.2 mol/l sodium phosphate buffer containing 10 mmol/l EDTA (stock buffer, pH 7.2), 100 µl of 6 mmol/l 5,5'-dithiobis-(2-nitrobenzoic acid) in the stock buffer, and 190 µl of water were added into the microfuge tube. Ten microliters of 250 iu/ml glutathione reductase were added with mixing to initiate the reaction. The formation of 5-thio-2-nitrobenzoic acid was detected photometrically by a change of absorption at 412 nm. The total GSH content is calculated according to a standard curve. The total amount of GSH calculated was divided by the number of oocytes in the sample to obtain the content per oocyte (pmol/oocyte).

In vitro fertilization

COCs cultured for a total of 44 h in maturation medium were stripped of cumulus mechanically by gentle aspiration with a pipette. Oocytes were washed three times with TALP medium and groups of 30–35 oocytes were transferred into each well of a four-well multi-dish containing 250 µl IVF medium previously equilibrated at 38.5°C under 5% CO₂. The sperm-rich fraction of semen from a mature, fertility-tested boar was collected by the gloved-hand method and immediately transported to the laboratory diluted 1:8 in Beltsville thawing solution. The semen samples were centrifuged (1200 g, 3 min) and the resultant sperm pellets were diluted in TALP medium at the desired concentration (Matás *et al.* 2003). The sperm suspension (250 µl) was added to each fertilization well to obtain a final concentration of 3000 cells/oocyte. At 5 h post-insemination (hpi), oocytes were washed twice with fresh NCSU-23 by gentle aspiration through a glass pipette and allowed to continue in culture at 38.5°C under 5% CO₂ until fixation or transfer.

At 18 hpi, a sample of oocytes was stained with Hoechst 33342 as described above and examined at × 400 magnification for evidence of sperm penetration and pronuclear formation under an epifluorescence microscope.

Embryo transfer

Multiparous (two pregnancies) crossbred sows with synchronized estrus cycles were used for surgical embryo transfer. Estrus was checked daily in the presence of a mature teaser boar. Occurrence of estrus was defined by the standing reflex in front of a boar (back-pressure test) and reddening and swelling of the vulva. Oviducts from animals which had showed heat 72 h before embryo transfer were exposed through mid-ventral incision under general anesthesia, as previously described (Coy *et al.* 1993). Over 100 potential embryos (22–24 hpi) were introduced in one oviduct by a Tom Cat Catheter (Kendall Co., Mansfield, MA, USA) connected to an insulin syringe containing PBS at 37°C. The catheter was later observed under a stereomicroscope to check that all embryos had been transferred into the oviducts. The sows were kept under the usual farm conditions, and 25–28 days after transfer pregnancy diagnosis was carried out by ultrasonography (Echoscanner T-100).

Experimental design





The experimental design is schematically represented in Fig. 1.

Before running the experiment, 831 oocytes in four replicates were stained just after collection with BCB to verify the growth stage and check if culture in roscovitine for 22 h inhibited the last step of growing. The numbers of oocytes examined were 264, 301 and 266 for before culture group, ROS group and A group respectively (see below for details).

Experiment 1. Nuclear status after 22 h culture in roscovitine

This experiment was carried out to establish whether culture of oocytes in NCSU-37 medium without dibutyryl cAMP, eCG and hCG and with 50 µmol/l roscovitine for the first 22 h of maturation (ROS group, *n* = 152) kept the nuclear stage at the same level as found in oocytes just after recovery (Before culture group, *n* = 156). As control groups, oocytes cultured for 22 h in supplemented NCSU-37 medium were used under usual IVM conditions, which included dibutyryl cAMP, eCG and hCG (A group, *n* = 173) and oocytes cultured for 22 h in the same NCSU-37 medium without dibutyryl cAMP, eCG and hCG (B group, *n* = 164). Therefore, it could be seen whether the inhibitory effect of roscovitine on the nuclear progression was similar, or dissimilar, to that demonstrated by dibutyryl cAMP, which only blocks oocytes at stages before the GV-II stage (Funahashi *et al.* 1997); the nuclear stage of inhibited oocytes (ROS and A groups) at 22 h and of those under supposed spontaneous resumption of meiosis (B group) could also be compared.

After culture, oocyte nuclear status (GV-0 to Met II) was recorded in all groups. This experiment was performed in four replicates.

Experiment 1. Nuclear Status After 22 h Culture in 50 μ M Roscovitine**Experiment 2. Nuclear Progression After IVM With a Prematuration Period in Roscovitine****Experiment 3. Oocyte Glutathione Content After Roscovitine Treatment****Experiment 4. In Vitro Fertilization and Full Development of Roscovitine Treated Oocytes****Figure 1** Experimental design. ET: Embryo transfer.**Experiment 2. Nuclear progression after IVM with a pre-maturation period in roscovitine**

In order to assess the reversibility of the treatment with roscovitine, COCs were cultured for 44 h under permissive maturation conditions in each of three replicates with (ROS-IVM group, $n = 153$) or without (IVM group, $n = 161$) a previous culture period of 22 h in the presence of 50 μ mol/l roscovitine. Nuclear stage was recorded by Hoechst staining.

Experiment 3. Oocyte GSH content after roscovitine treatment

This experiment was designed to examine the influence of roscovitine on the intracellular GSH content of oocytes just after collection, at 22 h of culture with or without roscovitine, and after maturation with or without a pre-culture in roscovitine. In seven replicates, 960 COCs were collected and allotted to five groups: (1) Before culture group (oocytes processed just after collection); (2) A group (oocytes cultured in supplemented NCSU-37 medium with dibutyl cAMP, eCG and hCG); (3) ROS group (oocytes cultured for 22 h in supplemented NCSU-37 medium without dibutyl cAMP, eCG and hCG and with

50 μ mol/l roscovitine); (4) IVM group (oocytes cultured for 44 h under the described step-wise IVM system); (5) ROS-IVM group (oocytes cultured for 22 h as the ROS group and for a further 44 h as the IVM group). GSH content was measured as described above.

Experiment 4. IVF and full development of roscovitine-treated oocytes

Just after recovering, 631 oocytes (in four replicates) were cultured in the presence of 50 μ mol/l roscovitine for 22 h. Following this period of time, oocytes were washed and allowed to mature for 44 h under permissive conditions (ROS-IVM group). Other 610 COCs were also collected and introduced in the IVM system at the same time as the ROS-IVM group. Matured oocytes from both groups were then fertilized and 18 hpi, samples of the potential embryos ($n = 133$ and $n = 127$ for ROS-IVM and IVM groups respectively) were processed to assess penetration and monospermy rates. The remaining cells (498 for ROS-IVM and 483 for ROS groups) were transferred into the oviduct of physiologically synchronized sows (one sow per group and replicate, eight sows in total). Pregnancies were assessed by ultrasonography 25–28 days after transfer.

Statistical analysis

Data are presented as means \pm S.E.M. and all percentages were modeled according to the binomial model of variables. The variables in all the experiments were analyzed by one-way ANOVA. These variables were as follows: experiments 1 and 2, the nuclear stage (GV-0 to Met II); experiment 3, the GSH content; experiment 4, the percentage of penetration, the number of sperm cells per penetrated oocyte, the male pronuclear formation and the putative embryos (putative embryos were defined as the percentage of monospermic oocytes with two pronuclei with respect to the total number of penetrated oocytes). When ANOVA revealed a significant effect, values were compared using Tukey's test. *P* values < 0.05 were taken to denote statistical significance.

Results

After collection, $86.0 \pm 2.1\%$ of oocytes showed a blue cytoplasm coloration following BCB staining and were classified as grown oocytes. After 22 h of culture in medium with or without roscovitine, this proportion increased significantly to 95.0 ± 1.2 and $97.0 \pm 1.0\%$ respectively.

Nuclear status after 22 h in roscovitine

Roscovitine was effective in inhibiting the meiotic resumption after 22 h of culture, giving a higher percentage of GV-I stage oocytes and a lower percentage of GV-III stage oocytes in the ROS group than in A or B groups (Table 1). Just before culture and after treatment with roscovitine, oocytes were similar for both nuclear stages. The percentage of oocytes reaching the Met-I stage was higher in the B group than in the remaining three groups.

Nuclear progression after IVM with a 22-h pre-maturation period in roscovitine

After maturation for 44 h, oocytes from the ROS-IVM group resumed meiosis and reached Met II in the same proportions as in the IVM group, over 85% (Table 2).

Oocyte GSH content after roscovitine treatment

As shown in Fig. 2, intracellular GSH content increased after oocytes were cultured. When oocytes were kept for 22 h in culture, GSH content increased from 2.24 to 10.74 pmol/oocyte, and still higher when roscovitine was present during culture (15.23 pmol/oocyte). After IVM, no differences were observed independently of a 22 h pre-maturation period with (ROS-IVM) or without (IVM) roscovitine, being 7.02 and 7.48 pmol/oocyte respectively. However, both groups showed a lower amount of GSH than the ROS group.

IVF and full development of roscovitine-treated oocytes

No differences were observed in IVF results after insemination of oocytes treated (ROS-IVM) or not (IVM) with roscovitine. Total data from four replicates showed a penetration rate of 73.0 ± 3.9 and $78.7 \pm 3.6\%$ respectively for ROS-IVM and IVM groups, and from the penetrated oocytes 41.2 ± 5.0 and $37.0 \pm 4.8\%$ respectively were monospermic. Male pronuclear formation was >87% in both groups.

After embryo transfer, both ROS-IVM and IVM sows from the second replicate became pregnant and delivered 12 (four male/eight female) and ten (five male/five female) healthy piglets respectively (Table 3). The weight of animals (means \pm S.E.M.), assessed 42 days after birth, was 8.3 ± 0.5 and 11.5 ± 0.6 kg respectively for ROS-IVM and IVM groups. Sows from replicates 1, 3 and 4 showed heat estrus 21–24 days after embryo transfer.

Discussion

A principal disadvantage of slaughterhouse material is the heterogeneity among batches of ovaries that could be responsible for variable results. In the present study, the growing stage of the oocytes was first assessed in four trials by BCB staining demonstrating that the selection of the follicles by slicing was appropriate, with the proportion of fully grown oocytes always >80%.

Table 1 Nuclear status of pig oocytes after 22 h culture in NCSU-37 with or without hormonal supplements, or with 50 μ mol/l roscovitine.

Group	Before culture	A	B	ROS	P value
<i>n</i>	156	173	164	152	
GV-0	96 ± 2.4^a	0.6 ± 0.6^b	0^b	0^b	<0.001
GV-I	60.9 ± 3.9^a	37.0 ± 3.7^b	31.7 ± 3.6^b	63.2 ± 3.9^a	<0.001
GV-II	8.3 ± 2.2^a	37.0 ± 3.7^b	26.2 ± 3.5^b	29.6 ± 3.7^b	<0.001
GV-III	7.0 ± 2.1^a	23.7 ± 3.2^b	20.1 ± 3.1^b	7.2 ± 2.1^a	<0.001
GV-IV	7.0 ± 2.1^a	0^b	3.7 ± 1.5^{ab}	0^b	<0.001
Metaphase I	5.1 ± 1.8^{ab}	1.2 ± 0.8^a	13.4 ± 2.7^b	0^a	<0.001
Anaphase I	0	0	1.2 ± 0.9	0	0.12
Telophase I	0.6 ± 0.6	0	2.4 ± 1.2	0	0.06
Metaphase II	1.3 ± 0.9	0.6 ± 0.6	2.4 ± 1.2	0	0.53

Different superscript letters in the same row indicate that values are significantly different ($P < 0.05$). A; 22 h culture in NCSU-37 with hCG, eCG and dibutyryl cAMP. B; 22 h culture in NCSU-37 without hCG, eCG and dibutyryl camp. ROS; 22 h culture in NCSU-37 without hCG, eCG and dibutyryl cAMP, and with 50 μ mol/l roscovitine.

Table 2 Nuclear status of pig oocytes after *in vitro* maturation with (ROS-IVM) or without (IVM) a prematuration period of 22 h in 50 $\mu\text{mol/l}$ roscovitine.

Group	IVM	ROS-IVM	P value
n	161	153	
GV-0	0.6 \pm 0.6	0	0.33
GV-I	6.2 \pm 1.9	4.6 \pm 1.7	0.52
GV-II	1.9 \pm 1.1	0	0.09
GV-III	1.2 \pm 0.9	1.3 \pm 0.9	0.96
GV-IV	0 ^a	6.5 \pm 2.0 ^b	<0.001
Metaphase I	1.9 \pm 1.1	0.6 \pm 0.6	0.34
Anaphase I	0	0	1
Telophase I	0.6 \pm 0.6	1.3 \pm 0.9	0.53
Metaphase II	87.6 \pm 2.6	85.6 \pm 2.8	0.61

Different superscript letters in the same row indicate that values are significantly different ($P < 0.05$).

Interestingly, the inhibitory effect of roscovitine on oocyte growth (measured as the activity of glucose-6-phosphate dehydrogenase) was not observed, since the proportion of BCB+ oocytes increased up to 95% after 22 h of culture. This could indicate that the two-step IVM system is useful for producing more developmentally competent oocytes. Although there are slight indications of a possible beneficial effect of pre-culture in roscovitine on further embryo development (Mermillod *et al.* 2000, Marchal *et al.* 2001, Coy *et al.* 2003, 2004), interpretation of these results must be cautious since an influence of roscovitine on the inhibition of RNA synthesis has also been demonstrated (Ljungman & Paulsen 2001). Moreover, recent studies have shown that roscovitine did not prevent most of the modifications of the protein phosphorylation pattern observed during maturation (Vigneron *et al.* 2004a) and that some pathways involved in the regulation of bovine oocyte maturation seem to be independent of MPF activity




Figure 2 Intracellular GSH content in pig oocytes before culture, after 22 h culture in NCSU-37 with hCG, eCG and dibutyryl cAMP. A group; without hCG, eCG, dibutyryl cAMP. ROS group; without hCG, eCG, dibutyryl cAMP and with 50 μM roscovitine. IVM group; after a conventional IVM (22 h A + 22 h B). ROS-IVM group; after conventional IVM (22 h A + 22 h B) with preculture in 50 $\mu\text{mol/l}$ roscovitine. Different letters indicate significant differences ($P < 0.001$).

and meiotic resumption (Vigneron *et al.* 2004b). However, delaying meiotic resumption (without any aim of improving competence) may be of interest from a practical point of view for any of the mentioned reasons.

The effect of roscovitine on the resumption of meiosis has been shown previously in pigs. Ju *et al.* (2003) found that levels of 80–120 $\mu\text{mol/l}$ roscovitine (Sigma, R-7772) were necessary to inhibit GVBD in 83–91% of oocytes. However, they assessed the nuclear stage of oocytes at 44 h from the beginning of culture in roscovitine, when the activity of the inhibitor could have been decreased in the medium (Meijer & Raymond 2003). In our study, doses of 50 $\mu\text{mol/l}$ roscovitine were enough to block the GVBD in >90% of oocytes after 22 h of culture with similarities between control and ROS groups being evident for the GV-I and GV-III stages. In contrast, when the IVM medium without roscovitine was employed, oocytes started to resume meiosis spontaneously at this time. Moreover, when the NCSU-37 medium without dibutyryl cAMP was used, the progression to stages beyond GV-4 was faster, since the inhibitory effect of dibutyryl cAMP (Funahashi *et al.* 1997) was absent. These data first demonstrate that progression of porcine oocytes to the GV-III stage in the presence of 50 $\mu\text{mol/l}$ roscovitine is inhibited in a different manner than in the presence of dibutyryl cAMP or in the absence of inhibitors, since they remain at similar nuclear stages to those in the follicles (as assessed just after recovery).

The reversibility of the roscovitine effect on the nuclear maturation has been corroborated in this study by the high proportions of Met II stage oocytes reached after 44 h of culture in the conventional IVM system following the 22 h of inhibition. However, it is well documented that nuclear and cytoplasmic maturation are not always correlated (Coy *et al.* 1999). The resumption of meiosis in oocytes after removal from roscovitine, presumably re-starting the activation of MPF, is not a prerequisite to ensure the cytoplasmic maturation of such oocytes. On the contrary, intracellular GSH content and the ability of oocytes to form a male pronucleus are commonly accepted parameters for assessing cytoplasmic maturation of porcine oocytes (Funahashi *et al.* 1995, Sawai *et al.* 1997, Coy *et al.* 1999). GSH is produced via the γ -glutamyl cycle and is dependent on the amount of cysteine available to the cell (Meister & Tate 1976). Cysteine is commonly added to pig oocyte maturation medium as a substrate for GSH synthesis to promote male pronuclear formation (Yoshida *et al.* 1993) at a 0.57 mmol/l concentration, and this was the case for NCSU-37. In this study, the GSH content of immature oocytes just after removal from follicles was lower than that found after 22 or 44 h of culture in IVM medium, the latter value being at the same baseline level as reported by Brad *et al.* (2003). The IVM system employed was presumably effective for oocyte GSH synthesis.

The GSH content was higher at 22 h of culture than at 44 h, in contrast to the findings of Yoshida *et al.* (1993)

Table 3 IVF results and pregnancies after surgical transfer of porcine embryo derived from *in vitro*-matured oocytes with (ROS-IVM) or without (IVM) a prematuration period of 22 h in 50 μ mol/l roscovitine.

Group	Replicate	N*	Penetration (%)	Spermatozoa/oocyte	Male pronucleus (%)	Putative embryos** (%)	Number Transferred***	Pregnancy	Piglets born
IVM	1	30	90.0 \pm 5.6	2.51 \pm 0.3	100	33.3 \pm 9.2	110	–	
ROS-IVM	1	25	60.0 \pm 10.0	1.53 \pm 0.2	86.7 \pm 9.1	40.0 \pm 13.1	110	–	
IVM	2	19	47.4 \pm 11.8	1.56 \pm 0.2	88.9 \pm 11.1	44.4 \pm 17.6	123	+	10
ROS-IVM	2	30	63.3 \pm 8.9	1.47 \pm 0.2	84.2 \pm 8.6	57.9 \pm 11.6	117	+	12
IVM	3	37	86.5 \pm 5.7	2.53 \pm 0.3	100	34.4 \pm 8.5	110	–	
ROS-IVM	3	46	84.8 \pm 5.3	2.54 \pm 0.2	87.2 \pm 5.4	23.1 \pm 6.8	150	–	
IVM	4	41	78.0 \pm 6.5	2.09 \pm 0.3	87.1 \pm 6.1	40.6 \pm 8.8	140	–	
ROS-IVM	4	32	75.0 \pm 7.8	1.54 \pm 0.2	95.8 \pm 4.2	58.3 \pm 10.3	121	–	
ANOVA									
Roscovitine			0.38	0.05	0.20	0.38			
Replicate			0.0015	0.0018	0.73	0.06			
Roscv \times repl.			0.48	0.24	0.13	0.37			

* Number of stained embryos 18 post-insemination.

** Proportion of zygotes with two visible pronuclei from the penetrated oocytes.

*** Number of transferred embryos 22–24 h post-insemination.

who have reported a continuous increase in GSH content during IVM. On the one hand, these authors employed TCM-199 for IVM, which is a rich culture medium with readily available GSH precursors (cysteine, cystine, glutamic acid, glutamine and glycine) and even glutathione. However, in our study NCSU-37 was used and was supplemented only with cysteine and glutamine (glutamine is the first cystine pre-cursor auto-oxidized, within 1 h, in the maturation medium (de Matos & Furnus 2000) that cannot be incorporated so easily as cysteine (Yoshida & Takahashi 1998)). On the other hand, it has been reported that GSH synthesis during initial and mid phases of porcine oocyte maturation is related to the acquisition of sperm nuclear decondensing ability (Yoshida 1993) and GSH synthesis during the initial stages of hamster oocyte maturation is responsible for the higher GSH levels in mature oocytes (Perreault *et al.* 1988). These observations suggest a strong GSH synthesis during the first half of oocyte maturation and would be in agreement with our results showing a GSH peak after 22 h of culture.

The present study also shows that oocytes pre-cultured in roscovitine for 22 h had a higher level of GSH than those cultured with dibutyryl cAMP for the same time. Cystine uptake activity is abolished by the mechanical disruption of gap junctional communication (Yoshida & Takahashi 1998), and GSH synthesis may be impaired due to the uncoupling of cumulus cells (de Matos *et al.* 1997). Since roscovitine-treated oocytes do not show cumulus expansion as shown by dibutyryl cAMP-treated oocytes (Marchal *et al.* 2001, Coy *et al.* 2004, and personal observations in the present experiments), the closest cooperation between cumulus cells and oocyte would be maintained for a longer time, explaining in part the higher GSH content observed in roscovitine-treated oocytes. Also, the inhibitory effect of roscovitine must be considered not only on the cdk but also on RNA synthesis (Ljungman & Paulsen 2001). Because GSH is involved in multiple functions, including DNA and protein synthesis

(Lafleur *et al.* 1994), it could be expected that oocytes treated with roscovitine showed a higher GSH content than their counterparts, as was observed in our case since this functional activity of GSH was probably decreased.

Regarding male pronuclear formation, our data first show that roscovitine does not affect this parameter. This result was expected due to the similar levels of GSH found for *in vitro*-matured oocytes pre-cultured or not with roscovitine, since it has been reported that male pronuclear formation is related to intracellular GSH content (Yoshida *et al.* 1992, Funahashi *et al.* 1995). Moreover, no differences were found for any of the fertilization parameters assessed at 18 hpi, in agreement with most previous studies in bovine (Mermillod *et al.* 2000, Ponderato *et al.* 2001, Coy *et al.* 2005).

Finally, the key result in our study was the pregnancy of one recipient sow and the delivery of 12 healthy piglets when roscovitine-pre-cultured embryos were transferred. It is well documented that the *in vitro* production of pig embryos with the ability to develop to term is difficult due to polyspermy and deficient embryo culture media (Coy & Romar 2002, Funahashi 2003). Only 20–30% of transferred embryos survive despite considerable improvement in the techniques for *in vitro* production of porcine embryos (Abeydeera 2002), and approximately six piglets per sow after surgical embryo transfer are obtained (Abeydeera *et al.* 1998, Kikuchi *et al.* 1999). With these premises, and taking into account that about 37 and 41% of the transferred zygotes in IVM and ROS-IVM could be expected to develop to term, a high number of putative embryos was transferred at only 22–24 h of insemination and *in vitro* culture, in order to ensure a sufficient number of viable embryos in the oviduct of the synchronized sow. Thus, a final success of 25% for both IVM and ROS-IVM groups could be considered satisfactory and is among the normal rates reported in the literature.

In conclusion, this study demonstrates that pig oocytes cultured for 22 h with 50 μ mol/l roscovitine remained at

similar nuclear stages to oocytes just after recovery. Cytoplasmic maturation assessed as intracellular GSH content and male pronuclear formation were not affected detrimentally by culture with roscovitine. Moreover, embryos obtained from this two-step IVM system can be developed to term showing the final competence of porcine oocytes pre-cultured with roscovitine, thus making its introduction desirable in assisted reproductive technologies programs.

Acknowledgements

The authors would like to thank: Professor RHF Hunter for critical reading of the manuscript and help with the English; Manuel Sansegundo, David Gumbao, Jose Guillermo Velasquez and Diana Patricia Barajas for their generous technical assistance with embryo transfer; Lo Navarro Farm for providing the boar semen and the animals necessary for this study; and the staff of the slaughterhouse El Pozo for supplying the ovaries. This research was supported by Ministerio de Ciencia y Tecnología, Spain (AGL2003-03144).

References

- Abeysdeera LR 2002 In vitro production of embryos in swine. *Theriogenology* **57** 256–273.
- Abeysdeera LR, Wang WH, Cantley TC, Rieke A, Prather RS & Day BN 1998 Presence of epidermal growth factor during in vitro maturation of pig oocytes and embryo culture can modulate blastocyst development after in vitro fertilization. *Molecular Reproduction and Development* **51** 395–401.
- Anderson ME 1985 Determination of glutathione and glutathione disulfide in biological samples. In *Methods in Enzymology: Glutamate, Glutamine, Glutathione and Related Compounds*, vol. 113, pp 548–555. Ed A Meister. New York: Academic Press.
- Avery B, Hay-Schmidt A, Hyttel P & Greve T 1998 Embryo development, oocyte morphology, and kinetics of meiotic maturation in bovine oocytes exposed to 6-dimethylaminopurine prior to in vitro maturation. *Molecular Reproduction and Development* **50** 334–344.
- Brad AM, Bormann CL, Swain JE, Durkin RE, Johnson AE, Clifford AL & Krisher RL 2003 Glutathione and adenosine triphosphate content of in vivo and in vitro matured porcine oocytes. *Molecular Reproduction and Development* **64** 492–498.
- Coy P & Romar R 2002 In vitro production of pig embryos: a point of view. *Reproduction Fertility and Development* **14** 275–286.
- Coy P, Martínez E, Ruiz S, Vázquez JM, Roca J, Matas C & Pellicer MT 1993 In vitro fertilization of pig oocytes after different incubation intervals. *Theriogenology* **39** 1201–1208.
- Coy P, Ruiz S, Romar R, Campos I & Gadea J 1999 Maturation, fertilization and complete development of porcine oocytes matured under different systems. *Theriogenology* **51** 799–812.
- Coy P, Gadea J, Romar R, Matás C & García E 2002 Effect of in vitro fertilization medium on the acrosome reaction, cortical reaction, zona pellucida hardening, and in vitro development in pigs. *Reproduction* **124** 279–288.
- Coy P, Cánovas S, García E, Ruiz S, Gadea J, Matás C & Romar R 2003 Effect of roscovitine, a cyclin-dependent kinases inhibitor, on pig in vitro maturation and fertilization by ICSI. *Human Reproduction* **18** 37.
- Coy P, Cánovas S, Ruiz S, Matás C, Gadea J, García E, Sansegundo M, García F, Gumbao D & Romar R 2004 First pregnancies after transferring of in vitro produced pig embryos pre-cultured with roscovitine. *Human Reproduction* **19** 41–42.
- Coy P, Romar R, Payton RR, McCann L, Saxton AM & Edwards JL 2005 Maintenance of meiotic arrest in bovine oocytes using the S-enantiomer of roscovitine: effects on maturation, fertilization and subsequent embryo development *in vitro*. *Reproduction* **129** 19–26.
- de Matos DG & Furnus CC 2000 The importance of having high glutathione (GSH) level after bovine *in vitro* maturation on embryo development effect of beta-mercaptoethanol, cysteine and cystine. *Theriogenology* **53** 761–771.
- de Matos DG, Furnus CC & Moses DF 1997 Glutathione synthesis during in vitro maturation of bovine oocytes: role of the cumulus cells. *Biology of Reproduction* **57** 1420–1425.
- Dode MAN & Adona PR 2001 Developmental capacity of *Bos indicus* oocytes after inhibition of meiotic resumption by 6-dimethylaminopurine. *Theriogenology* **65** 171–180.
- Faerge I, Mayes M, Hyttel P & Sirard MA 2001 Nuclear ultrastructure in bovine oocytes after inhibition of meiosis by chemical and biological inhibitors. *Molecular Reproduction and Development* **59** 459–467.
- Fair T, Hyttel P, Motlik J, Boland M & Lonergan P 2002 Maintenance of meiotic arrest in bovine oocytes *in vitro* using butyrolactone I: effects on oocyte ultrastructure and nucleolus function. *Molecular Reproduction and Development* **62** 375–386.
- Funahashi H 2003 Polyspermic penetration in porcine IVM-IVF systems. *Reproduction, Fertility and Development* **15** 167–177.
- Funahashi H & Day BN 1993 Effects of the duration of exposure to supplemental hormones on cytoplasmic maturation of pig oocytes *in vitro*. *Journal of Reproduction and Fertility* **98** 179–185.
- Funahashi H, Stumpf TT, Cantley TC, Kim NH & Day BN 1995 Pronuclear formation and intracellular glutathione content of in vitro-matured porcine oocytes following in vitro fertilization and/or electrical activation. *Zygote* **3** 273–281.
- Funahashi H, Cantley TC & Day BN 1997 Synchronization of meiosis in porcine oocytes by exposure to dibutyryl cyclic AMP improves developmental competence following in vitro fertilization. *Biology of Reproduction* **57** 49–53.
- Gibbons J, Arat S, Rzucidlo J, Miyoshi K, Waltenburg R, Respass D, Venable A & Stice S 2002 Enhanced survivability of cloned calves derived from roscovitine-treated adult somatic cells. *Biology of Reproduction* **66** 895–900.
- Ju J-C, Tsay C & Ruan C-W 2003 Alterations and reversibility in the chromatin, cytoskeleton and development of pig oocytes treated with roscovitine. *Molecular Reproduction and Development* **64** 482–491.
- Kikuchi K, Kashiwazaki N, Noguchi J, Shimada A, Takahashi R, Hirabayashi M, Shino M, Ueda M & Kaneko H 1999 Developmental competence, after transfer to recipients, of porcine oocytes matured, fertilized, and cultured *in vitro*. *Biology of Reproduction* **60** 336–340.
- Kitagawa M, Okabe T, Ogino H, Matsumoto H, Suzuki-Takahashi I, Kokubo T, Higashi H, Saitoh S, Taya Y & Yasuda H 1993 Butyrolactone I, a selective inhibitor of cdk2 and cdc2 kinase. *Oncogene* **8** 2425–2432.
- Krischek C & Meinecke B 2001 Roscovitine, a specific inhibitor of cyclin-dependent protein kinases, reversibly inhibits chromatin condensation during in vitro maturation of porcine oocytes. *Zygote* **9** 309–316.
- Kubelka M, Motlik J, Schultz RM & Pavlok A 2000 Butyrolactone I reversibly inhibits meiotic maturation of bovine oocytes, without influencing chromosome condensation activity. *Biology of Reproduction* **62** 292–302.
- Lafleur MV, Hoorweg JJ, Joenje H, Westmijze EJ & Retel J 1994 The ambivalent role of glutathione in the protection of DNA against singlet oxygen. *Free Radical Research* **21** 9–17.
- Liu L, Ju JC & Yang X 1998 Differential inactivation of maturation-promoting factor and mitogen-activated protein kinase following parthenogenetic activation of bovine oocytes. *Biology of Reproduction* **59** 537–545.

- Ljungman M & Paulsen MT** 2001 The cyclin-dependent kinase inhibitor roscovitine inhibits RNA synthesis and triggers nuclear accumulation of p53 that is unmodified at Ser15 and Lys382. *Molecular Pharmacology* **60** 785–789.
- Lonergan P, Khatir H, Carolan C & Mermillod P** 1997 Bovine blastocyst production *in vitro* after inhibition of oocyte meiotic resumption for 24 h. *Journal of Reproduction and Fertility* **109** 355–365.
- Machaty Z, Day BN & Prather RS** 1998 Development of early porcine embryos *in vitro* and *in vivo*. *Biology of Reproduction* **59** 451–455.
- Marchal R, Tomanek M, Terqui M & Mermillod P** 2001 Effects of cell cycle dependent kinases inhibitor on nuclear and cytoplasmic maturation of porcine oocytes. *Molecular Reproduction and Development* **60** 65–73.
- Matás C, Coy P, Romar R, Marco M, Gadea J & Ruiz S** 2003 Effect of sperm preparation method on *in vitro* fertilisation in pigs. *Reproduction* **125** 133–141.
- Meijer L & Kim SH** 1997 Chemical inhibitors of cyclin-dependent kinases. *Methods in Enzymology* **283** 113–128.
- Meijer L & Raymond E** 2003 Roscovitine and other purines as kinase inhibitors. From starfish to clinical trials. *Accounts of Chemical Research* **36** 417–425.
- Meister A & Tate SS** 1976 Glutathione and related gamma-glutamyl compounds: biosynthesis and utilization. *Annual Reviews of Biochemistry* **45** 559–604.
- Mermillod P, Tomanek M, Marchal R & Meijer L** 2000 High developmental competence of cattle oocytes maintained at the germinal vesicle stage for 24 hours in culture by specific inhibition of MPF kinase activity. *Molecular Reproduction and Development* **55** 89–95.
- Motlik J, Pavlok A, Kubelka M, Kalous J & Kalab P** 1998 Interplay between CDC2 kinase and MAP kinase pathway during maturation of mammalian oocytes. *Theriogenology* **49** 461–419.
- Motlik J, Pavlok A, Lapathitis G & Kubelka M** 2000 Impact of two-step *in vitro* culture systems on developmental potency of oocytes. *Reproduction in Domestic Animals* **35** 267–271.
- Payton R, Romar R, Coy P, Saxton AM, Lawrence JL & Edwards JL** 2004 Susceptibility of bovine germinal vesicle-stage oocytes from antral follicles to direct effects of heat stress *in vitro*. *Biology of Reproduction* **71** 1303–1308.
- Perreault SD, Barbee RR & Slott VL** 1988 Importance of glutathione in the acquisition and maintenance of sperm nuclear decondensing activity in maturing hamster oocytes. *Developmental Biology* **125** 181–186.
- Petters RM & Wells KD** 1993 Culture of pig embryos. *Journal of Reproduction and Fertility Supplement* **48** 61–73.
- Picton HM & Gosden RG** 1999 Oogenesis, in mammals. In *Encyclopedia of Reproduction*, Vol 3, pp 488–497. New York: Academic Press.
- Ponderato N, Lagutina I, Crotti G, Turín P, Galli C & Lazzari G** 2001 Bovine oocytes treated prior to *in vitro* maturation with a combination of butyrolactone I and roscovitine at low doses maintain a normal developmental capacity. *Molecular Reproduction and Development* **60** 579–685.
- Ponderato N, Crotti G, Turini P, Duchi R, Galli C & Lazzari G** 2002 Embryonic and foetal development of bovine oocytes treated with a combination of butyrolactone I and roscovitine in an enriched medium prior to IVM and IVF. *Molecular Reproduction and Development* **62** 513–518.
- Rath D, Long CR, Dobrinsky JR, Welch GR, Schreiber LL & Johnson LA** 1999 *In vitro* production of sexed embryos for gender preselection: high-speed sorting of X-chromosome bearing sperm to produce pigs after embryo transfer. *Journal of Animal Science* **77** 3346–3352.
- Roca J, Martínez E, Vazquez JM & Lucas X** 1998 Selection of immature pig oocytes for homologous *in vitro* penetration assays with the brilliant cresyl blue test. *Reproduction, Fertility and Development* **10** 479–485.
- Rodríguez-González E, Lopez-Bejar M, Velilla E & Paramio MT** 2002 Selection of prepubertal goat oocytes using the brilliant cresyl blue test. *Theriogenology* **57** 1397–1409.
- Sawai K, Funahashi H & Niwa K** 1997 Stage-specific requirement of cysteine during *in vitro* maturation of porcine oocytes for glutathione synthesis associated with male pronuclear formation. *Biology of Reproduction* **57** 1–6.
- Sirard MA, Richard F & Mayes M** 1998 Controlling meiotic resumption in bovine oocytes: a review. *Theriogenology* **49** 483–497.
- Vignerón C, Perreau C, Dalbiès-Tran R, Joly C, Humblot P, Uzbekova S & Mermillod P** 2004a Protein synthesis and mRNA storage in cattle oocytes maintained under meiotic block by roscovitine inhibition of MPF activity. *Molecular Reproduction and Development* **69** 457–465.
- Vignerón C, Perreau C, Dupont J, Uzbekova S, Prigent C & Mermillod P** 2004b Several signaling pathways are involved in the control of cattle oocyte maturation. *Molecular Reproduction and Development* **69** 466–474.
- Wu B, Ignatz GG, Currie WB & Yang X** 1996 Temporal distinctions in the synthesis and accumulation of proteins by oocytes and cumulus cells during maturation *in vitro* of bovine oocytes. *Molecular Reproduction and Development* **45** 560–565.
- Yoshida M** 1993 Role of glutathione in the maturation and fertilization of pig oocytes *in vitro*. *Molecular Reproduction and Development* **35** 76–81.
- Yoshida M & Takahashi K** 1998 Changes in glutathione and transport activity of cysteine and cystine during maturation, fertilization, and development in pig ova. In *Gametes: Development and Functions*, abstract 621. Eds A Lauria, F Gandolfi, G Enne, L Gianaroli. Rome: Sero Symposium.
- Yoshida M, Ishigaki K & Pursel VG** 1992 Effect of maturation media on male pronucleus formation in pig oocytes matured *in vitro*. *Molecular Reproduction and Development* **31** 68–71.
- Yoshida M, Ishigaki K, Nagai T, Chikyu M & Pursel VG** 1993 Glutathione concentration during maturation and after fertilization in pig oocytes: relevance to the ability of oocytes to form male pronucleus. *Biology of Reproduction* **49** 89–94.

Received 14 February 2005

First decision 4 March 2005

Revised manuscript received 22 March 2005

Accepted 24 March 2005