

Bifidobacterium mitigates autoimmune hepatitis by regulating IL-33-induced Treg/Th17 imbalance via the TLR2/4 signaling pathway

Jianguo Song^{1,2*}, Juan Dai^{2*}, Xueping Chen^{4*}, Fei Ding², Yanbo Ding², Liang Ma^{1,2} and Liwen Zhang³

¹Department of Gastroenterology, The Fifth People's Hospital of Xinjiang Uygur Autonomous Region, Xin Jiang, ²Department of Gastroenterology, The First People's Hospital of Changzhou, The Third Affiliated Hospital of Soochow University, ³Department of Pediatrics, the Second People's Hospital of Changzhou, Affiliated Hospital of Nanjing Medical University, Changzhou, Jiangsu and ⁴Department of Gastroenterology, The People's Hospital of Wujia, Xinjiang, China

*These authors made equal contributions to this study

Summary. The present work aims to evaluate the efficacy of Live Combined *Bifidobacterium*, *Lactobacillus* and *Enterococcus* Capsules (LCBLECs), a probiotic drug containing *Bifidobacterium*, in the treatment of autoimmune hepatitis (AIH). In this study, a mouse model of experimental autoimmune hepatitis (EAH) was established to investigate the effects of LCBLECs on AIH. The results showed that LCBLECs improved dysbiosis of gut microbiota, reduced liver injury, restored liver function, and maintained Treg/Th17 balance in EAH mice. In addition, LCBLECs restored Treg/Th17 balance in EAH mice by downregulating IL-33 production. Besides, LCBLECs also suppress IL-33 upregulation in EAH mice by inhibiting the TLR2/4 signaling pathway. Furthermore, LCBLECs also mitigated dysbiosis of gut microbiota and enhanced the efficacy of conventional treatment for AIH patients. To sum up, our findings revealed that LCBLECs exerted therapeutic effects on EAH mice by improving Treg/Th17 imbalance in an IL-33-dependent manner via the TLR2/4 signaling pathway and relieved the clinical symptoms of AIH patients, indicating *Bifidobacterium* supplementation with LCBLECs might be a potential adjuvant therapy for AIH treatment.

Key words: *Bifidobacterium*, Autoimmune hepatitis, IL-33, Treg/Th17 imbalance, TLR2/4

Corresponding Author: Liang Ma, Department of Gastroenterology, The First People's Hospital of Changzhou, The Third Affiliated Hospital of Soochow University, Changzhou, Jiangsu, 213000, People's Republic of China. e-mail: maliang212@126.com or Liwen Zhang, Department of Pediatrics, the Second People's Hospital of Changzhou, Affiliate Hospital of Nanjing Medical University, Changzhou, Jiangsu, China, 213000, People's Republic of China. e-mail: zhangliwenlove827@126.com

www.hh.um.es. DOI: 10.14670/HH-18-669

Introduction

Autoimmune hepatitis (AIH) is a chronic inflammatory liver disease characterized by interface hepatitis, circulating autoantibodies, and hypergammaglobulinemia (Yuksel et al., 2015). If left untreated, AIH may develop rapidly into liver cirrhosis and failure (Durazzo et al., 2019). Although AIH etiology and pathogenesis remain largely unknown, it is generally believed that AIH usually occurs in genetically susceptible hosts as a consequence of excessive immune responses towards liver tissue (van Gerven et al., 2016). Regulatory T cells (Tregs) play a crucial role in the maintenance of immune balance to prevent autoimmune diseases (Scheinecker et al., 2020). As a member of the T-helper effector cell family, interleukin IL-17-producing Th cells (Th17s) can induce inflammation and autoimmune tissue injury by promoting production of proinflammatory cytokines (Singh et al., 2013). Studies have reported that Treg/Th17 imbalance contributes to liver injury in the pathological process of liver diseases (Zhang et al., 2019; Lu et al., 2021; Zhao et al., 2022). Recently, it has been demonstrated that Treg/Th17 imbalance is associated with poor prognosis of AIH patients (Vuerich et al., 2021). However, the underlying mechanism of Treg/Th17 imbalance in AIH remains to be further understood.

As key organs in nutrient absorption and metabolism, the gut and the liver are closely connected through the biliary tract, portal vein and systemic circulation (Zeuzem, 2000; Nie et al., 2021). This bidirectional interaction between gut and the liver is called the gut-liver axis (Song and Zhang, 2022). As evidenced by previous studies, the alteration of gut microbiota composition plays a crucial role in the cross-talk of the gut-liver axis and is closely associated with inflammation, oxidative stress, and lipid deposition in



the liver (Meng et al., 2018). Studies have also shown that the gut-liver axis exists and plays a crucial role in the development of liver diseases, including AIH (Wang et al., 2022b). It is widely recognized that probiotics are beneficial regulators of gut microbiota, among which *Bifidobacterium* is a probiotic that has been widely studied and utilized to inhibit the growth of pathogenic bacteria in the intestinal tract, maintain intestinal health, and protect the intestinal barrier (Wang et al., 2022a). In addition, prophylactic administration of *Bifidobacterium* also has a favorable impact on AIH (Zhang et al., 2020). Although it has been reported that *Bifidobacterium* can improve immune function by regulating Th17/Treg balance (Dong et al., 2022; Sapra et al., 2022), it is still unclear whether *Bifidobacterium* can protect against AIH by improving Th17/Treg imbalance.

The present study aimed to investigate the efficacy and the specific mechanisms of *Bifidobacterium* in AIH treatment using Live Combined *Bifidobacterium*, *Lactobacillus* and *Enterococcus* Capsules (LCBLECs), a probiotic mix containing *Bifidobacterium*. Based on the results of LCBLEC administration for experimental autoimmune hepatitis (EAH) mice and AIH patients, it was found that LCBLECs alleviated the symptoms of EAH mice and promoted the efficacy of conventional treatment for AIH patients, suggesting that oral supplementation of *Bifidobacterium* might exhibit potential therapeutic effects on AIH.

Materials and methods

Patients

20 AIH patients at active phase were selected from AIH patients diagnosed in The Third Affiliated Hospital of Soochow University. AIH diagnosis was based on Simplified Criteria for the Diagnosis of Autoimmune Hepatitis promulgated by International Autoimmune Hepatitis Group (IAIHG) in 2008 and Practice Guidelines for Diagnosis and Management of Autoimmune Hepatitis promulgated by American Association for the Study of Liver Disease (AASLD) in 2010. Criteria for AIH patients at active phase: 1) fatigue, arthralgia, jaundice; 2) serum alanine aminotransferase (ALT) level >10-fold of upper limit of normal (ULN); 3) serum aspartate aminotransferase (AST) level \geq 5-fold of ULN, and γ -globulin level \geq 2-fold of ULN; 4) histological findings of bridging necrosis or multilobular necrosis. Each patient enrolled provided written informed consent. The study was approved by the Ethics Committee of The Third Affiliated Hospital of Soochow University.

The enrolled patients were randomly assigned to Group I (conventional treatment; n=10) and Group II (conventional treatment + LCBLECs; n=10). All AIH patients in Group II were given LCBLECs (0.66g \times 3/d t.i.d.). 2 weeks later, serum samples and fecal samples were collected from AIH patients in Group I and Group II for further analysis.

Animal study

Wildtype (WT) and IL-33 knockout (IL-33^{-/-}) female C57BL/6 mice (6-8 weeks old; 18-20g) were purchased from Model Animal Research Center of Nanjing University and kept in specific pathogen-free conditions. Mice were randomly divided into six groups (n=6): Control (WT), EAH (WT), EAH (WT)+LCBLECs, Control (IL-33^{-/-}), EAH (IL-33^{-/-}), and EAH (IL-33^{-/-})+LCBLECs. To induce EAH, mice received intraperitoneal injection of S-100 liver antigen emulsified in complete Freund's adjuvant (CFA) on Day 0 and Day 7, as previously described (Liang et al., 2021). From day 15 to day 28, LCBLECs were administered by gavage (0.345 g/kg/day) (Zhao et al., 2013). Finally, all mice were sacrificed to collect serum samples, liver tissues and feces for further analysis. All animal procedures were permitted by Ethics Committee of The Third Affiliated Hospital of Soochow University (Approval No.: [2019]KY048-01).

Hematoxylin and eosin (H&E) staining

H&E staining was applied for histopathological evaluation of liver tissues. In brief, liver tissues were fixed in 4% paraformaldehyde for 48h, embedded in paraffin, prepared into sections (4 μ m), deparaffinized in xylene, and rehydrated. Then, the sections were examined under light microscopy for tissue injury, necrosis, and infiltrating leukocytes.

Biochemical analysis

ALT, aspartate aminotransferase (AST), Immunoglobulin G (IgG), Immunoglobulin M (IgM), and Immunoglobulin A (IgA) levels in serum samples from mice and AIH patients were determined using the automated chemistry analyzer (BioMajesty, Japan).

Table 1. Primers used for RT-qPCR analysis of gene expression.

Gene	Primer sequence 5'-3'
IL-33	Forward: TTCCTGTCTGTATTGAGAAACCT Reverse: TTTGCCGGGGAAATCTTGA
sST2	Forward: CCAGCCCTTCATCTGGGTATC Reverse: ATGGTGTGTTCACTAGGCGG
TLR2	Forward: ATCCCCTTCCCCTCACTCCA Reverse: GCCCGGAGCCTAGGAGT
TLR4	Forward: TCCCTGCATAGAGGTAGTCCC Reverse: TCAAGGGGTTGAAGCTCAGA
MyD88	Forward: GATGACCCCTAGGACAAACG Reverse: ACTGATATCGTTGGGGCAG
IRAK1	Forward: CCTCAGATCCCAGGAACAGGTAT Reverse: AGTTCTCCTGTGGGGAAGGG
TRAF6	Forward: GGAGTTTGACCCACCTCTGG Reverse: CTTGTGCCCTGCATCCCTTA

Bifidobacterium mitigates autoimmune hepatitis

RT-qPCR

To detect abundance of gut microbiota in fecal samples, bacterial genomic DNA was extracted from fecal samples and purified for RT-qPCR analysis. The abundances of *Lactobacillus*, *Bifidobacterium*, *Clostridium leptum*, and *Bacteroides fragilis*, *Escherichia coli* were quantified by RT-qPCR in StepOne™ Real-Time PCR System using SYBR® Premix Ex Taq™ reagents.

To detect gene expression, total RNA was extracted from liver tissues of mice using TRIzol (Invitrogen, CA) and reversely transcribed into cDNA using Quantitect Reverse Transcription Kit (Qiagen, USA). Then, RT-qPCR was performed in StepOne™ Real-Time PCR System using SYBR® Premix Ex Taq™ reagents. Relative gene expression was normalized to β -actin and quantified by $2^{-\Delta\Delta Ct}$ method. The primers used for RT-qPCR were as follows in Table 1.

ELISA

The concentrations of IL-33 and its soluble receptor ST2 (sST2) in serum samples from AIH patients and liver tissue supernatant from mice were determined using IL-33/sST2 ELISA kit (R&D Systems, USA) as per manufacturer's instructions.

Gas chromatography-mass spectrometry (GC-MS)

The concentrations of short-chain fatty acids (SCFAs) in fecal samples from mice and AIH patients

were determined by GC-MS as previously described (Jiang et al., 2020).

Flow cytometry

Flow cytometry was applied to detect Th17s and Tregs in liver tissues. Briefly, liver tissues were aseptically removed from mice and prepared into monocyte suspensions. After intracellular staining with specific anti-cytokine antibodies, the number of Th17s (CD3⁺CD4⁺IL-17a⁺) and Tregs (CD3⁺CD4⁺CD25⁺FOXP3⁺) was determined by FACS flow cytometry. The antibodies used for flow cytometry analysis were as follows: anti-CD3 (ab11089; Abcam), anti-CD4 (ab133616; Abcam), anti-IL-17a (ab302922; Abcam), anti-CD25 (Cat. No.: 561780; R&D Systems), and anti-FOXP3 (ab215206; Abcam).


Western blotting

Total protein was isolated from liver tissue by RIPA Buffer (Beyotime), separated by 10% SDS/PAGE and electroblotted onto PVDF membranes. Next, PVDF membranes were blocked with 5% skimmed milk. Then, the membranes were incubated with primary antibody at 4°C overnight and treated with secondary antibody at 37°C for 45 min. Finally, the protein bands were visualized using BeyoECL Plus (Beyotime). The protein level of sST2 was detected by using an anti-ST2 antibody. Primary antibodies used for western blotting assay were: anti-IL-33 antibody (ab187060; Abcam), anti-ST2 antibody (ab228543), anti-TLR2 antibody


A

Gut microbiota	Control (WT)	EAH (WT)	EAH+LCBLECs (WT)
Lactobacillus	1±0.125	0.313±0.103	0.642±0.105
Bifidobacterium	1±0.161	0.269±0.213	0.651±0.113
Clostridium leptum	1±0.201	0.356±0.113	0.613±0.121
Bacteroides fragilis	1±0.134	0.312±0.104	0.689±0.211
Escherichia coli	1±0.113	2.131±0.211	1.377±0.131

B



C



D




Fig. 1. LCBLECs ameliorate dysbiosis of gut microbiota in EAH mice. **A.** Lactobacillus, *Bifidobacterium*, *Clostridium leptum*, *Bacteroides fragilis*, and *Escherichia coli* levels in feces of mice from Control (WT) (n=5), EAH (WT) (n=5), and EAH (WT)+LCBLECs (n=5) groups by RT-qPCR. **B.** acetic acid, butyric acid, and propionic acid levels in feces of mice by GC-MS. * $P < 0.05$; ** $P < 0.01$.

(ab213676; Abcam), anti-TLR4 antibody (Cat. No.: 48-2300; Invitrogen), anti-MyD88 antibody (ab219413; Abcam), anti-IRAK1 antibody (ab238; Abcam), anti-TRAF6 antibody (ab33915; Abcam), anti-NF- κ B p-p65 antibody (ab76302; Abcam), anti-NF- κ B p65 antibody (ab32536; Abcam), and anti-GAPDH antibody (ab8245; Abcam). Secondary antibodies used were: Anti-Rabbit/Mouse horseradish peroxidase (HRP)-conjugated secondary antibodies (ab97085, ab98799; Abcam).

Statistical analysis

All data were expressed as mean \pm SD. Comparisons were performed by one-way Analysis of Variance (ANOVA) using SPSS 19.0 (SPSS Inc., USA). Graphs were plotted using GraphPad Prism 6.0. Values with $P < 0.05$ were deemed significant in statistics.

Results

LCBLECs ameliorate dysbiosis of gut microbiota in EAH mice

EAH mice exhibited a significant decrease in *Lactobacillus*, *Bifidobacterium*, *Clostridium leptum*, and *Bacteroides fragilis* levels and a significant increase in *Escherichia coli* level compared with mice in the control group, which was substantially abated by LCBLEC administration (Fig. 1A). As important metabolites of gut microbiota, SCFAs, including acetic acid, butyric acid, and propionic acid, are deeply involved in immune regulation and autoimmunity by regulating the generation of Th17s and Tregs (Haase et al., 2018). Therefore, acetic acid, butyric acid, and propionic acid levels in feces of mice from each group were measured.




Fig. 2. LCBLECs alleviate EAH severity in mice. **A.** HE staining of liver tissues of mice from Control (WT) (n=5), EAH (WT) (n=5), and EAH (WT)+LCBLECs (n=5) groups. **B-F.** serum levels of liver enzymes (ALT and AST) and autoantibodies (IgG, IgA, and IgM) by biochemical analysis. * $P < 0.05$; ** $P < 0.01$

Bifidobacterium mitigates autoimmune hepatitis

As indicated in Fig. 1B-D, LCBLECs significantly eliminate the reduction of acetic acid, butyric acid, and propionic acid levels in EAH mice. Therefore, LCBLECs effectively mitigated EAH-induced dysbiosis of gut microbiota.

LCBLECs alleviate EAH severity in mice

To explore the effect of LCBLECs on AIH progression, we established an EAH model based on C57BL/6 mice. H&E staining results showed that LCBLECs significantly attenuated EAH-induced liver injury, including destruction of liver structure, large areas of necrosis, and infiltration of inflammatory cells, in mice (Fig. 2A). In addition, serum levels of liver enzymes (ALT and AST) and autoantibodies (IgG, IgA, and IgM) were also determined. As shown in Fig. 2B-F, LCBLECs partially reversed the elevation of ALT, AST, IgG, IgA, and IgM levels in serum samples of EAH mice. The above results demonstrated that LCBLECs exerted liver-protective effects on EAH mice.

LCBLECs relieve Treg/Th17 imbalance in EAH mice

It has been demonstrated that *Bifidobacterium* could

restore Treg/Th17 imbalance in autoimmune diseases (Lopez et al., 2016). To explore the specific effects of LCBLECs on Treg/Th17 balance in EAH mice, flow




Fig. 3. LCBLECs relieve Treg/Th17 imbalance in EAH mice. **A, B.** Detection of the frequency of Treg (CD3⁺CD4⁺CD25⁺FOXP3⁺) cells and Th17 (CD3⁺CD4⁺IL-17a⁺) cells by flow cytometry. **P*<0.05; ***P*<0.01.




Fig. 4. LCBLECs regulate Treg/Th17 imbalance in an IL-33-dependent manner. **A, B.** IL-33 and sST2 mRNA and protein levels in liver tissues of mice from Control (WT) (n=5), EAH (WT) (n=5), and EAH (WT)+LCBLECs groups (n=5) by RT-qPCR and Western blotting. **C.** Serum IL-33 and sST2 levels by ELISA. **D, E.** Detection of the frequency Treg (CD3⁺CD4⁺CD25⁺FOXP3⁺) cells and Th17 (CD3⁺CD4⁺IL-17a⁺) cells in liver tissues of mice from Control (WT) (n=5), EAH (WT) (n=5), EAH (WT)+LCBLECs (n=5), Control (IL-33^{-/-}) (n=5), EAH (IL-33^{-/-}) (n=5), and EAH (IL-33^{-/-})+LCBLECs (n=5) groups by flow cytometry. **P*<0.05; ***P*<0.01.

cytometry was applied to detect the frequencies of Tregs and Th17s in liver tissues of mice. The results showed that LCBLEC treatment markedly increased the percentages of Tregs and reduced the number of Th17s in liver tissues from EAH mice (Fig. 3A,B). These results indicated that LCBLECs might help restore Treg/Th17 balance in liver tissues from EAH mice.


LCBLECs regulate Treg/Th17 imbalance in an IL-33-dependent manner

As a new member of the IL-1 family, IL-33 plays a critical role in AIH pathogenesis and severity (Abe et al., 2019). In addition, IL-33 has also been identified as a key factor for maintenance of Treg/Th17 balance in numerous diseases (Cheng et al., 2022). Hence, it was hypothesized that LCBLECs might regulate Treg/Th17 balance in EAH mice via modulating IL-33 secretion. It was shown that LCBLECs inhibited IL-33 and sST2 levels in liver tissues (Fig. 4A,B) and serum samples (Fig. 4C) from EAH mice. Next, we used IL-33^{-/-} mice to further investigate the role of IL-33 in the restorative effects of LCBLECs on EAH-induced Treg/Th17 imbalance. Treatment with LCBLECs reversed EAH-induced reduction of Tregs and increase of Th17s in liver tissues from WT mice, while IL-33^{-/-} mice showed

few changes in the frequencies of Tregs and Th17s (Fig. 4D,E). Therefore, LCBLECs relieved Treg/Th17 imbalance EAH mice via IL-33.

LCBLECs suppress EAH-induced IL-33 upregulation by deactivating the TLR2/4 signaling pathway

Toll-like receptors (TLRs), a group of pattern-recognition receptors (PRRs), are essential regulators of both innate and adaptive immune responses (Yu et al., 2017). Among them, TLR2 and TLR4 have been identified as key regulators of signaling pathways in AIH (Cai et al., 2022; Zhou et al., 2022). *Bifidobacterium* has been identified as an inhibitor of both TLR2 and TLR4 signaling pathway (Al-Sadi et al., 2021; Chen et al., 2021). Therefore, the expression of related genes and proteins in TLR2/TLR4 signaling pathway were detected. As shown in Fig. 5A, TLR2 and TLR4, MyD88, IRAK1 and TRAF6 mRNA expressions significantly increased in liver tissues from EAH mice, which was effectively abrogated by treatment with LCBLECs. Furthermore, Western blotting results indicated that LCBLECs exerted inhibitory effects on TLR2, TLR4, MyD88, IRAK1, TRAF6, and NF- κ B p-p65 protein levels in EAH mice (Fig. 5B). Taken together, LCBLECs might repress IL-33 upregulation in



Bifidobacterium mitigates autoimmune hepatitis

EAH mice by inhibiting the hyperactivation of the TLR2/4 signaling pathway.

LCBLECs improve clinical symptoms of AIH patients

To determine the clinical effect of LCBLECs on AIH, AIH patients were randomly assigned to Group I (conventional treatment; n=10) and Group II (conventional treatment + LCBLECs; n=10). AIH patients in Group II were orally administrated with LCBLECs (0.66g×3/d t.i.d.). As shown in Fig. 6A, AIH patients in Group II exhibited higher *Lactobacillus*, *Bifidobacterium*, *Clostridium leptum*, and *Bacteroides fragilis* levels and lower *Escherichia coli* level than AIH patients in Group I. Similarly, LCBLECs further increased acetic acid, butyric acid, and propionic acid levels in feces from AIH patients (Fig. 6B). Biochemical results revealed that LCBLEC administration further promoted the reduction of ALT, AST, IgG, IgA, and IgM levels in AIH patients induced by conventional treatment (Fig. 6C-G). Moreover, LCBLEC treatment further decreased serum IL-33 and sST2 levels of AIH patients, compared with conventional treatment (Fig. 6H,I). Taken together, cotreatment with LCBLECs exhibited better effects on clinical symptoms of AIH patients than conventional treatment alone.

Discussion

AIH incidence has been rising in recent years, which

is a significant threat to human health globally (Feld and Heathcote, 2003). Therefore, it is necessary to find novel and effective therapeutic agents for AIH treatment. Probiotics exert beneficial effects in the prevention and treatment of autoimmune disorders by restoring altered gut microbiota, maintaining intestinal barrier, and regulating immune homeostasis (Shahbazi et al., 2020). As a major genus of intestinal probiotics, *Bifidobacterium* has been studied for its immunoregulatory role in autoimmune diseases (Rinaldi et al., 2019). However, there were only few studies on the application of *Bifidobacterium* in AIH treatment. In this work, LCBLECs substantially improved EAH-induced dysbiosis of gut microbiota in mice, as evidenced by the increase in *Lactobacillus*, *Bifidobacterium*, *Clostridium leptum*, and *Bacteroides fragilis* population and the decrease in *Escherichia coli* population. In addition, LCBLECs reduced ALT, AST, IgG, IgA, and IgM levels in serum samples from EAH mice, indicating LCBLECs might alleviate EAH severity by reducing liver injury and restoring liver function. Furthermore, LCBLECs also restored the altered gut microbiota in AIH patients and significantly improved the efficacy of conventional treatment for AIH patients. The above results suggested that LCBLECs altered the composition of gut microbiota, thereby exerting liver-protective effects in AIH.

Previous studies have reported the crucial role of the Treg/Th17 balance in autoimmune diseases (Lee, 2018). Our team has also previously reported the importance of




Fig. 6. LCBLECs improve clinical symptoms of AIH patients. **A.** *Lactobacillus*, *Bifidobacterium*, *Clostridium leptum*, *Bacteroides fragilis*, and *Escherichia coli* levels in feces of AIH patients from Group I (n=10) and Group II (n=10). **B.** acetic acid, butyric acid, and propionic acid levels in feces from AIH patients. **C-G.** ALT, AST, IgG, IgA, and IgM levels in serum samples from AIH patients. **H, I.** IL-33 and sST2 levels in serum samples from AIH patients. * $P<0.05$; ** $P<0.01$.

the Treg/Th17 balance in AIH pathogenesis (Liang et al., 2018). In addition, *Bifidobacterium* has been proven as an important regulator of Treg/Th17 balance in autoimmune diseases (Shi et al., 2018). Therefore, we explored how *Bifidobacterium* affect Treg/Th17 balance in AIH progression. The results showed that LCBLECs reversed the reduction of Tregs and the increase of Th17s in liver tissues from EAH mice. These results indicated that LCBLECs may exert therapeutic effects on AIH by maintaining the Treg/Th17 balance. As documented in previous studies, IL-33 plays an important role in AIH pathogenesis and severity (Abe et al., 2019). In addition, IL-33 also serves as an inducer for Treg/Th17 imbalance in LPS-stimulated acute respiratory distress syndrome (Cheng et al., 2022). Hence, we assumed that LCBLECs might restore Treg/Th17 balance in EAH mice via inhibiting IL-33 production. Our results confirmed that administration with LCBLECs effectively abated EAH-induced increase of IL-33 and sST2 levels. Besides, LCBLEC treatment further promoted the decrease in serum IL-33 and sST2 levels of AIH patients caused by conventional treatment. Furthermore, LCBLEC treatment significantly increased the frequencies of Tregs and reduced the frequencies of Th17s in EAH-induced WT mice; however, the frequencies of Tregs and Th17s in IL-33^{-/-} mice had barely changed, implying LCBLECs regulated the balance between Tregs and Th17s in EAH mice via IL-33.

TLRs play critical roles in the innate recognition of pathogens besides bridging both innate and adaptive immune responses (Kawai and Akira, 2011). Among them, TLR2 and TLR4 have been identified as key regulators of autoimmune response in AIH (Chi et al., 2018). Therefore, we wondered whether TLR2 or TLR4 signaling was involved in LCBLEC-mediated downregulation of IL-33 in EAH mice. Herein, LCBLECs decreased TLR2, TLR4, MyD88, IRAK1 and TRAF6 mRNA expression in liver tissues from EAH mice. In addition, TLR2, TLR4, MyD88, IRAK1, TRAF6, and NF- κ B p-p65 protein levels also declined after treatment with LCBLECs. These results indicated that LCBLECs might reinstate the Treg/Th17 balance in EAH mice by regulating IL-33 production via the TLR2/4 signaling pathway.

The EAH model used in this study is a chronic EAH model which has been widely applied in the exploration of the potential mechanisms of AIH (Ma et al., 2007). As an acute EAH model, Concanavalin A (ConA)-induced EAH model has also been frequently used in the investigation of T-cell function in AIH (Zheng et al., 2018). In this study, we have demonstrated the protective role of LCBLECs in a chronic EAH model. Therefore, we can speculate that LCBLECs may also exert therapeutic effects in ConA-induced EAH, which will be further verified in future studies.

Conclusion

To sum up, our findings revealed that LCBLECs

exerted therapeutic effects on EAH mice by improving dysbiosis of gut microbiota and Treg/Th17 balance. In addition, it was also demonstrated that LCBLECs restored Treg/Th17 balance in EAH mice in an IL-33-dependent manner via the TLR2/4 signaling pathway. Furthermore, LCBLECs also mitigated dysbiosis of gut microbiota and enhanced the efficacy of conventional treatment for AIH patients. Hence, *Bifidobacterium* supplementation with LCBLECs might be a potential adjuvant therapy for AIH treatment.

Funding. This study was sponsored by grants from the Top Talent of Changzhou "The 14th Five-Year Plan" High-Level Health Talents Training Project (2022260), Changzhou Medical Center, Nanjing Medical University (CMCB202301); Changzhou Municipal Health Commission Science and Technology Program (No. QN202316; No. QN202342 and No. QN202343); Natural Science Foundation of Xinjiang Uygur Autonomous Region (2021D01F64); Natural Science Foundation of Xinjiang Uygur Autonomous Region (2022D01A307); Xinjiang Kirgiz Autonomous Prefecture Medical and Health Technology Project (Grant No. 43; No.44); the National Natural Science Foundation of China (No. 81700500).

References

- Abe K., Takahashi A., Fujita M., Hayashi M., Okai K., Nozawa Y. and Ohira H. (2019). Interleukin-33/st2-mediated inflammation plays a critical role in the pathogenesis and severity of type I autoimmune hepatitis. *Hepatology*. 3, 670-684.
- Al-Sadi R., Dharmaparakash V., Nighot P., Guo S., Nighot M., Do T. and Ma T.Y. (2021). *Bifidobacterium bifidum* enhances the intestinal epithelial tight junction barrier and protects against intestinal inflammation by targeting the toll-like receptor-2 pathway in an NF- κ B-independent manner. *Int. J. Mol. Sci.* 22, 8070.
- Cai T., Xu L., Xia D., Zhu L., Lin Y., Yu S., Zhu K., Wang X., Pan C., Chen Y. and Chen D. (2022). Polyguanine alleviated autoimmune hepatitis through regulation of macrophage receptor with collagenous structure and TLR4-TRIF-NF- κ B signalling. *J. Cell. Mol. Med.* 26, 5690-5701.
- Chen Y., Yang B., Stanton C., Ross R.P., Zhao J., Zhang H. and Chen W. (2021). *Bifidobacterium pseudocatenulatum* ameliorates dss-induced colitis by maintaining intestinal mechanical barrier, blocking proinflammatory cytokines, inhibiting tlr4/nf-kappab signaling, and altering gut microbiota. *J. Agric. Food Chem.* 69, 1496-1512.
- Cheng L., Jiao Y., Jiang W., Zhang X., Zhang L. and Jia G. (2022). IL-33 deficiency attenuates lung inflammation by inducing Th17 response and impacting the Th17/Treg balance in LPS-induced ARDS mice via dendritic cells. *J. Immunol. Res.* 2022, 9543083.
- Chi G., Feng X.X., Ru Y.X., Xiong T., Gao Y., Wang H., Luo Z.L., Mo R., Guo F., He Y.P., Zhang G.M., Tian D.A. and Feng Z.H. (2018). TLR2/4 ligand-amplified liver inflammation promotes initiation of autoimmune hepatitis due to sustained IL-6/IL-12/IL-4/IL-25 expression. *Mol. Immunol.* 99, 171-181.
- Dong J., Ping L., Cao T., Sun L., Liu D., Wang S., Huo G. and Li B. (2022). Immunomodulatory effects of the *bifidobacterium longum* BL-10 on lipopolysaccharide-induced intestinal mucosal immune injury. *Front. Immunol.* 13, 947755.
- Durazzo M., Lupi G., Scandella M., Ferro A. and Gruden G. (2019).

Bifidobacterium mitigates autoimmune hepatitis

- Autoimmune hepatitis treatment in the elderly: A systematic review. *World J. Gastroenterol.* 25, 2809-2818.
- Feld J.J. and Heathcote E.J. (2003). Epidemiology of autoimmune liver disease. *J. Gastroenterol. Hepatol.* 18, 1118-1128.
- Haase S., Haghikia A., Wilck N., Muller D.N. and Linker R.A. (2018). Impacts of microbiome metabolites on immune regulation and autoimmunity. *Immunology* 154, 230-238.
- Jiang X.W., Li Y.T., Ye J.Z., Lv L.X., Yang L.Y., Bian X.Y., Wu W.R., Wu J.J., Shi D., Wang Q., Fang D.Q., Wang K.C., Wang Q.Q., Lu Y.M., Xie J.J. and Li L.J. (2020). New strain of *pediococcus pentosaceus* alleviates ethanol-induced liver injury by modulating the gut microbiota and short-chain fatty acid metabolism. *World J. Gastroenterol.* 26, 6224-6240.
- Kawai T. and Akira S. (2011). Toll-like receptors and their crosstalk with other innate receptors in infection and immunity. *Immunity* 34, 637-650.
- Lee G.R. (2018). The balance of Th17 versus treg cells in autoimmunity. *Int. J. Mol. Sci.* 19, 730.
- Liang M., Liwen Z., Yun Z., Yanbo D. and Jianping C. (2018). The imbalance between Foxp3⁺Tregs and Th1/Th17/Th22 cells in patients with newly diagnosed autoimmune hepatitis. *J. Immunol. Res.* 2018, 3753081.
- Liang M., Liwen Z., Jianguo S., Juan D., Fei D., Yin Z., Changping W. and Jianping C. (2021). Fecal microbiota transplantation controls progression of experimental autoimmune hepatitis in mice by modulating the TFR/TFH immune imbalance and intestinal microbiota composition. *Front. Immunol.* 12, 728723.
- Lopez P., de Paz B., Rodriguez-Carrio J., Hevia A., Sanchez B., Margolles A. and Suarez A. (2016). Th17 responses and natural IgM antibodies are related to gut microbiota composition in systemic lupus erythematosus patients. *Sci. Rep.* 6, 24072.
- Lu Z., Lu F., Wu L., He B., Chen Z. and Yan M. (2021). Berberine attenuates non-alcoholic steatohepatitis by regulating chemerin/CMKLR1 signalling pathway and Treg/Th17 ratio. *Naunyn Schmiedebergs Arch. Pharmacol.* 394, 383-390.
- Ma X., Jia Y.T. and Qiu D.K. (2007). Inhibition of p38 mitogen-activated protein kinase attenuates experimental autoimmune hepatitis: Involvement of nuclear factor kappa b. *World J. Gastroenterol.* 13, 4249-4254.
- Meng X., Li S., Li Y., Gan R.Y. and Li H.B. (2018). Gut microbiota's relationship with liver disease and role in hepatoprotection by dietary natural products and probiotics. *Nutrients* 10, 1457.
- Nie Y., Liu Q., Zhang W., Wan Y., Huang C. and Zhu X. (2021). Ursolic acid reverses liver fibrosis by inhibiting NOX4/NLRP3 inflammasome pathways and bacterial dysbiosis. *Gut Microbes* 13, 1972746.
- Rinaldi E., Consonni A., Cordiglieri C., Sacco G., Crasa C., Fontana A., Morelli L., Elli M., Mantegazza R. and Baggi F. (2019). Therapeutic effect of *bifidobacterium* administration on experimental autoimmune myasthenia gravis in lewis rats. *Front. Immunol.* 10, 2949.
- Sapra L., Shokeen N., Porwal K., Saini C., Bhardwaj A., Mathew M., Mishra P.K., Chattopadhyay N., Dar H.Y., Verma B. and Srivastava R.K. (2022). *Bifidobacterium longum* ameliorates ovariectomy-induced bone loss via enhancing anti-osteoclastogenic and immunomodulatory potential of regulatory B cells (Bregs). *Front. Immunol.* 13, 875788.
- Scheinecker C., Goschl L. and Bonelli M. (2020). Treg cells in health and autoimmune diseases: New insights from single cell analysis. *J. Autoimmun.* 110, 102376.
- Shahbazi R., Yasavoli-Sharahi H., Alsadi N., Ismail N. and Matar C. (2020). Probiotics in treatment of viral respiratory infections and neuroinflammatory disorders. *Molecules* 25, 4891.
- Shi P., Qu H., Nian D., Chen Y., Liu X., Li Q., Li Q., Wang C., Ye M. and Ma B. (2018). Treatment of guillain-barre syndrome with *Bifidobacterium infantis* through regulation of T helper cells subsets. *Int. Immunopharmacol.* 61, 290-296.
- Singh B., Schwartz J.A., Sandrock C., Bellemore S.M. and Nikoospour E. (2013). Modulation of autoimmune diseases by interleukin (IL)-17 producing regulatory T helper (Th17) cells. *Indian J. Med. Res.* 138, 591-594.
- Song Q. and Zhang X. (2022). The role of gut-liver axis in gut microbiome dysbiosis associated NAFLD and NAFLD-HCC. *Biomedicine* 10, 524.
- van Gerven N.M., de Boer Y.S., Mulder C.J., van Nieuwkerk C.M. and Bouma G. (2016). Auto immune hepatitis. *World J. Gastroenterol.* 22, 4651-4661.
- Vuerich M., Wang N., Kalbasi A., Graham J.J. and Longhi M.S. (2021). Dysfunctional immune regulation in autoimmune hepatitis: From pathogenesis to novel therapies. *Front. Immunol.* 12, 746436.
- Wang K., Chen Y., Cao J., Liang R., Qiao Y., Ding L., Yang X. and Yang Z. (2022a). Mechanism of huangqi-honghua combination regulating the gut microbiota to affect bile acid metabolism towards preventing cerebral ischaemia-reperfusion injury in rats. *Pharm. Biol.* 60, 2189-2199.
- Wang K., Wu W., Jiang X., Xia J., Lv L., Li S., Zhuge A., Wu Z., Wang Q., Wang S. and Li L. (2022b). Multi-omics analysis reveals the protection of gasdermin D in concanavalin A-induced autoimmune hepatitis. *Microbiol. Spectr.* 10, e0171722.
- Yu F., Zhang X., Tian S., Geng L., Xu W., Ma N., Wang M., Jia Y., Liu X., Ma J., Quan Y., Zhang C., Guo L., An W. and Liu D. (2017). Comprehensive investigation of cytokine- and immune-related gene variants in HBV-associated hepatocellular carcinoma patients. *Biosci. Rep.* 37, BSR20171263.
- Yuksel M., Wang Y., Tai N., Peng J., Guo J., Beland K., Lapiere P., David C., Alvarez F., Colle I., Yan H., Mieli-Vergani G., Vergani D., Ma Y. and Wen L. (2015). A novel "humanized mouse" model for autoimmune hepatitis and the association of gut microbiota with liver inflammation. *Hepatology* 62, 1536-1550.
- Zeuzem S. (2000). Gut-liver axis. *Int. J. Colorectal Dis.* 15, 59-82.
- Zhang H., Jiang Z. and Zhang L. (2019). Dual effect of t helper cell 17 (Th17) and regulatory T cell (Treg) in liver pathological process: From occurrence to end stage of disease. *Int. Immunopharmacol.* 69, 50-59.
- Zhang H., Liu M., Liu X., Zhong W., Li Y., Ran Y., Guo L., Chen X., Zhao J., Wang B. and Zhou L. (2020). *Bifidobacterium animalis* ssp. Lactis 420 mitigates autoimmune hepatitis through regulating intestinal barrier and liver immune cells. *Front. Immunol.* 11, 569104.
- Zhao H.M., Huang X.Y., Zuo Z.Q., Pan Q.H., Ao M.Y., Zhou F., Liu H.N., Liu Z.Y. and Liu D.Y. (2013). Probiotics increase T regulatory cells and reduce severity of experimental colitis in mice. *World J. Gastroenterol.* 19, 742-749.
- Zhao H., Guo P., Zuo Y., Wang Y., Zhao H., Lan T., Xue M., Zhang H. and Liang H. (2022). Folic acid intervention changes liver Foxp3 methylation and ameliorates the damage caused by Th17/Treg imbalance after long-term alcohol exposure. *Food Funct.* 13, 5262-5274.
- Zheng C., Yin S., Yang Y., Yu Y. and Xie X. (2018). Cd24 aggravates acute liver injury in autoimmune hepatitis by promoting IFN- γ

Bifidobacterium mitigates autoimmune hepatitis

production by CD4⁺ T cells. *Cell. Mol. Immunol.* 15, 260-271.
Zhou C., Zou Y., Zhang Y., Teng S. and Ye K. (2022). Involvement of
CCN1 protein and TLR2/4 signaling pathways in intestinal epithelial

cells response to *Listeria monocytogenes*. *Int. J. Mol. Sci.* 23, 2739.
Accepted October 18, 2023